

# CENTRAL COAST REGIONAL RESOURCE KIT

## METRIC DICTIONARY

Lead Contributors: Carol Clark, Peter Stine

Additional Contributions:

John Battles, Tadashi Moody, Patricia Manley, Michael Goulden, Liraz Bistriz, Joseph Stewart

9 May 2023, version 2.0

## TABLE OF CONTENTS

|  |    |
|--|----|
| Introduction                                       | 6  |
| What is the Regional Resource Kit effort?          | 6  |
| What this document is and its intended purpose     | 6  |
| Organizational Structure                           | 6  |
| Intended Purpose                                   | 8  |
| Fire Adapted Communities                           | 10 |
| Hazard   | 10 |
| Structure Exposure Score                           | 10 |
| Damage Potential                                   | 12 |
| Ember Load Index                                   | 14 |
| Ignition Cause                                     | 14 |
| Source of Ember Load to Buildings                  | 15 |
| Wildfire Hazard Potential                          | 16 |
| Fire Dynamics                                      | 17 |
| Functional Fire                                    | 17 |
| Fire Ignition Probability                          | 21 |
| Fire Return Interval Departure                     | 21 |
| Mean Percent FRI Departure, Since 1908             | 22 |
| Mean Percent FRI Departure, Since 1970             | 23 |
| FRID Condition Class for Departure                 | 23 |
| Time Since Last Fire                               | 24 |
| Current Fire Return Interval Departure, Since 1908 | 25 |
| Current Fire Return Interval Departure, Since 1970 | 25 |
| Severity   | 26 |
| Annual Burn Probability                            | 26 |
| Probability of High Fire Severity                  | 27 |
| Forest and Shrubland Resilience                    | 27 |
| Structure  | 28 |
| Density – Large Trees                              | 28 |
| Canopy Layer Count                                 | 29 |
| Canopy Veg Height                                  | 29 |
| Canopy Veg Cover                                   | 30 |
| Composition  | 31 |
| Tree Cover   | 31 |
| Shrub Cover  | 31 |
| Herbaceous Cover                                   | 32 |
| Seral Stage  | 32 |
| Disturbance  | 33 |
| Cumulative Tree Cover Loss                         | 33 |

|  |    |
|--|----|
| Cumulative Shrub Cover Lost  | 34 |
| Risk of Tree Dieoff During Drought   | 34 |
| Potential Climate Refugia -Baseline (Historical) Conditions  | 35 |
| Potential Climate Refugia - under Modeled Climate Change (MIROC model - hotter and drier)                                      | 36 |
| Potential Climate Refugia - Combined Modeled Climate Change (MIROC model - (Hotter and drier) and CNRM-CM5 (wetter and warmer) | 38 |
| Biodiversity Conservation  | 40 |
| Species Diversity  | 40 |
| Wildlife Species Richness  | 40 |
| Threatened/Endangered Vertebrate Species Richness  | 41 |
| Forest Raptors Species Richness  | 42 |
| Open habitat raptors species richness  | 43 |
| Hummingbirds species richness  | 43 |
| Focal Species  | 44 |
| California Spotted Owl   | 44 |
| Mountain Lion  | 45 |
| California red-legged frog   | 46 |
| loggerhead shrike  | 47 |
| Nuttall's woodpecker   | 47 |
| ringtail cat   | 48 |
| least bell's vireo   | 48 |
| Community Integrity  | 49 |
| Functional Group Species Richness  | 49 |
| Habitat Connectivity   | 50 |
| Economic Diversity   | 51 |
| Wood Product Industry  | 51 |
| Cost of Potential Treatments   | 51 |
| Carbon Sequestration   | 53 |
| Carbon Storage   | 53 |
| Total Aboveground Carbon   | 53 |
| Carbon Stability   | 54 |
| Aboveground Carbon Turnover Time   | 54 |
| Water Security   | 54 |
| Quantity   | 55 |
| Actual Evapotranspiration to Precipitation Fraction During Drought   | 55 |
| Precipitation Minus Actual Evapotranspiration during average conditions  | 56 |
| Quality  | 56 |
| Percent Impervious Surface   | 56 |
| Air Quality  | 57 |
| Particulate Matter   | 57 |

|   |    |
|---|----|
| Total Potential Smoke Emission - High Severity                      | 58 |
| Wetland Integrity   | 58 |
| Composition   | 58 |
| Aquatic Species Richness  | 59 |
| Wetland Diversity   | 59 |
| Riparian Habitat  | 60 |
| Social and Cultural Well-Being                                      | 60 |
| Environmental Justice   | 61 |
| Poverty Percentile  | 61 |
| Housing Burden Percentile   | 62 |
| Unemployment Percentile   | 63 |
| Operational Data Layers   | 64 |
| Fire  | 64 |
| Recent Fire Severity Class  | 64 |
| Housing Unit Density  | 65 |
| Wildland Urban Interface  | 66 |
| Administrative  | 67 |
| Urban-Agriculture Land Use  | 67 |
| Building Structure Density  | 67 |
| High-Use Recreation Areas   | 68 |
| Land Designations   | 68 |
| Ownership   | 69 |
| Roads   | 69 |
| Transmission Lines  | 70 |
| Distribution Lines  | 71 |
| Terrestrial   | 72 |
| CWHR Vegetation   | 72 |
| Aquatic   | 77 |
| Lakes/Reservoirs  | 77 |
| Perennial, Intermittent and Ephemeral Streams                       | 78 |
| Data Disclaimers  | 79 |
| Department of Forestry and Fire Protection (CALFIRE)                | 79 |
| California Department of Fish and Wildlife (CDFW)                   | 79 |
| Area of Conservation Emphasis (ACE)                                 | 79 |
| Biogeographic Information and Observation System (BIOS)             | 80 |
| Farmland Mapping and Monitoring Program (FMMP)                      | 81 |
| Open Street Map   | 81 |
| LANDFIRE  | 81 |
| California Forest Observatory (Salo Sciences)                       | 81 |
| Open Data Commons Open Database License (ODbL)                      | 81 |
| California Office of Environmental Health Hazard Assessment (OEHHA) | 82 |

|   |    |
|---|----|
| Center for Ecosystem Climate Solutions (CECS) – UC Irvine | 82 |
| Pyrologix   | 82 |
| USDA Forest Service (USFS)                                | 82 |
| US Fish and Wildlife Service (USFWS)                      | 82 |
| U.S. Geological survey (USGS)                             | 83 |
| Additional Resources                                      | 83 |
| References  | 83 |

# INTRODUCTION

## WHAT IS THE REGIONAL RESOURCE KIT EFFORT?

Reducing the risk of large, high intensity fire (and other mega-disturbances) through forest treatments has become a management imperative in California. A [Strategy for Shared Stewardship](#) (2018) and the USFS [Wildfire Crisis Implementation Plan](#) (2022) reinforce specific goals for pace and scale of strategic forest treatments over the next decade. Concurrently, the State of California has issued a new [Wildfire and Forest Resilience Action Plan](#) (2022), designed to strategically accelerate efforts to restore the health and resilience of California forests through a joint State of California - Forest Service framework to improve and enhance forest stewardship in California. The social incentives and the scientific knowledge to pursue meaningful restoration of forested landscapes in California are firmly established.

High quality geospatial data are an essential ingredient to address restoration/conservation of the broad suite of core socio-ecological values across landscapes, and to drive analytic tools for planning management investments. To support these initiatives an interagency team of scientists from the Forest Service/Pacific Southwest Research Station, California Natural Resources Agency/CALFIRE, and the University of California at Berkeley and University of California at Irvine collaborated on development of a comprehensive set of mapped data layers needed to accomplish large-scale landscape planning and restoration. Landscape level assessment using high quality data developed from ecological modeling techniques, informative analytical approaches and the resulting credible scientific outputs will be fundamental to inform and support large landscape restoration planning and execution.

The data layers included in this kit are meant to assist land managers in assessing their current landscape and plan for treatments to enhance resilience to human and natural disturbances. Thus each layer represents what the interagency team believes are the most relevant and reliable geospatial data available at this time. Each layer has been examined by the team and is supported by published data and/or was developed using standard methods. The methods for developing each layer are documented in the metric dictionary; however, the accuracy of each layer has not been quantified. It is anticipated that all data layers will be updated and refined as methods and source data evolve and improve.

## WHAT THIS DOCUMENT IS AND ITS INTENDED PURPOSE

---

### ORGANIZATIONAL STRUCTURE

This document has been organized to reflect the “Framework for Resilience” as set forth by the Tahoe Central Sierra Initiative (Manley et al. 2020, 2022). The framework is comprised of ten “**Pillars**” which support the full array of landscape management objectives that are inherently interdependent. Each pillar represents the desired long-term, landscape-scale outcome to restoring resilience. They include ecological values, such as biodiversity, as well as societal benefits to communities, such as water security. Within each pillar are “**Elements**” which represent the primary processes and core functions of that pillar, such as focal species, water quality, or economic health. Finally, within each element are the individual “**Metrics**” which describe the characteristics of elements in quantitative or qualitative terms. Metrics are used to assess, plan for, measure, and monitor progress toward desired outcomes and greater resilience.

The framework pillars are:

- **Fire Dynamics**
- **Forest Resilience**
- **Biodiversity Conservation**
- **Wetland Integrity**
- **Water Security**
- **Carbon Sequestration**
- **Air Quality**
- **Economic Diversity**
- **Fire Adapted Communities**
- **Social & Cultural Well-Being**

It is important to understand that while pillars and elements are consistent across California, the metrics used by a group may vary from region to region based on ecological and social differences (for example forest types or economy), available data, and the user preferences. It is equally important to recognize that due to the interdependent nature of the framework, some metrics overlap into multiple elements/pillars however have only been addressed a single time within this document.

The metrics are also divided into three "tiers." Among all these metrics, some are created and relevant statewide. Other metrics are more suited to conditions within a given region. The "Tiers" for metrics included in each RRK:

Tier 1 – metrics that are a single, consistent data layer, developed statewide; they can also be clipped to the boundary of the region so values within that region are the only ones included for calculations or regional statistics. Example: Annual Burn Probability.

Tier 2 – metrics relevant to a single region or relevant to multiple Regions but data layers differ among Regions because of varied data availability (sources) across Regions. Example: California gnatcatcher habitat suitability.

Tier 3 - metrics are those that would be of interest to some land managers for specific applications but not included as a core metric in an RRK. Example: Distribution of the Quino checkerspot butterfly.

Each RRK will contain all Tier 1 and Tier 2 data together to comprise the kit. Tier 3 data will be pointed to for reference and use, as needed.

**Within each Tier, the data layers are available in two forms: 1) data values native to the metric (raw), and 2) translated data values.** The raw data values are in the units of the metric, so for example the species richness map will show an estimated number of terrestrial vertebrate species per acre that can range from 0 to any number for each 30-m pixel, and the departure from historical fire return interval (FRID) map will have values that range from -100% to +100% departure. **The translated data values represent each metric using a common unit of measure with the same range of values from -1 to +1 that represent values that are generally considered favorable (+1) and unfavorable (-1).** In the case of species richness, higher species counts are considered more favorable and lower species counts are considered less favorable. In the case of FRID, values within the historical fire return interval are considered favorable, and high departure from the historical fire return interval is considered less favorable. In both cases, more and less favorable conditions for each metric are represented by values that range from +1 to -1 (respectively) so that multiple metrics can be evaluated together, including summarizing overall conditions at element and pillar levels to characterize socio-ecological resilience. The translated data are being developed and will be made available on the Central Coast RRK website as soon as they are ready.

Some data layers within this kit contain null values. We point this out here so users of the data will be aware and take whatever measures appropriate as they use and analyze the data. For some raster datasets in the RRK, areas have been masked (blanked) out and have a cell value of NoData (also referred to as null, NaN or missing). We, as producers and users of the data, cannot ignore NoData or fill them with zeros, since zero is often a valid value for some datasets. Removing NoData cells is not an option, a raster is a continuous grid. For users of the data performing further analyses and combining or "stacking" rasters, these NoData cells will mask out all values in that location in the output. To avoid this issue, the user must create values for the cells before combining them (i.e. 999 or any numeric value that is not real and clearly out of the range of the other values). Reasons for masking (blanking) out cells in RRK data:

- Cells are lakes or reservoirs
- Cells are urban or agriculture
- Cells contain no information relevant to the dataset (i.e. streams, habitat)
- Area (cells) subject to fire or other disturbance but the post disturbance condition or value is unknown

---

## INTENDED PURPOSE

Landscape level assessments, using high-quality data combined with decision support tools to help evaluate alternative treatment strategies, are fundamental to inform and support large landscape restoration planning. These data have been assembled in one place to provide comprehensive access for land managers.

Through this "metric dictionary," each metric has been defined to help end-users of the data (and for use with any decision support tools) to understand:

- What tier the metric is in (1, 2, or 3)
- Data vintage
- The definition meant by a given metric
- The expected use(s) of the metric
- The resolution of the developed data
- The data sources used to derive the metric
- The method of metric derivation
- The root file names

References have been included to help the reader understand potential methods for deriving metrics. It is our hope this information will help people make better use of all the assembled information and how it can best be used with various decision support tools. This dictionary will be updated periodically, as necessary.

Note that all metric data layers have been masked (i.e. blocked out) for open water (lakes, reservoirs) and a selected few have been masked for the urban and agricultural landscape (see the list of operational layers at the end of this document. This is done to avoid confusion with vegetation values coming from urban areas (e.g. city parks) or agricultural areas (e.g. irrigated farm land).

**The metrics (by Pillar) available in this Regional Resource Kit are listed below. The Tier 1 and Tier 2 metrics are included as part of the Kit. Tier 3 data are made available for applications, as needed, by the user. After each**



metric the source of the data is listed followed by a link to download a map of the data and to download the spatial data.

## FIRE ADAPTED COMMUNITIES

Wildfires are a keystone disturbance process in western US forests. However, the capacity for humans to coexist in the wildland urban interface (WUI) requires different restoration strategies aimed at the protection of life and property. This pillar evaluates the degree to which communities are living safely with fire and are accepting of management and natural ecological dynamics. It also evaluates the capacity for communities to manage desired, beneficial fire and suppress unwanted fire.

The definition of WUI used here, from Carlson et al 2022, adopts the definitions of interface and intermix WUI developed for previous census-based WUI mapping efforts based on U.S. Federal Register definitions (Radeloff et al., 2005; USDA & USDI, 2001). According to the definitions used for the building-based maps and for the census-based maps, WUI is where building density exceeds 6.17 units/km<sup>2</sup> and where land cover is either (1) at least 50% wildland vegetation (intermix) or (2) under 50% wildland vegetation but within 2.4 km (1.5 miles) of a patch of wildland vegetation at least 5 km<sup>2</sup> in area that contains at least 75% vegetation (interface). The distance selected for the interface definition is based on research from the California Fire Alliance suggesting that this is the average distance firebrands can travel from an active wildfire front (Stewart et al., 2007).

**DESIRED OUTCOME:** Communities have adapted to live safely in forested landscapes and understand the significance of fire to maintaining healthy forests. They have sufficient capacity to manage desired fire and suppress unwanted fire.

### HAZARD

The fire hazard element characterizes the fire risk in the wildland urban interface (WUI) defense and threat zones.

---

### STRUCTURE EXPOSURE SCORE

**Tier:** 1

**Data vintage:** The data are current through 2022.

**Metric Definition and Relevance:** This metric combines two data layers; one is the Wildland Urban Interface (WUI) as defined by Carlson et al. 2022 (see [WUI definition](#) section for more information), and a second data layer, Structure Exposure Score (SES), developed by Pyrologix LLC. The WUI includes the intermix and interface zones which collectively identify areas where structures occur and/or where structures are within a 1.5 miles wildland vegetation (see definition above) . The distance selected for the interface definition is based on research from the California Fire Alliance suggesting that this is the average distance firebrands can travel from an active wildfire front.

Structure Exposure Score is an integrated rating of wildfire hazard that includes the likelihood of a wildfire reaching a given location along with the potential intensity and ember load when that occurs. SES varies considerably across the landscape.

Pyrologix uses a standard geometric-interval classification to define the ten classes of SES, where each class break is 1.5 times larger than the previous break. So, homes located within Class X are 1.5 times more exposed than those in Class IX, and so on. This metric represents SES for WUI areas only.

- 1 (SES I): 0
- 2 (SES II): 0.01 to 50

3. 3 (SES III): 50 to 75
4. 4 (SES IV): 75 to 113
5. 5 (SES V): 113 to 169
6. 6 (SES VI): 169 to 253
7. 7 (SES VII): 253 to 380
8. 8 (SES VIII): 380 to 570
9. 9 (SES IX): 570 to 854
10. 10 (SES X): 854+

**Data Resolution:** 30m Raster

**Data Units:** Relative index, 10 classes

**Creation Method:**

**WUI:**

The current delineation of the WUI (Carlson et al. 2022) uses a mapping algorithm with definitions of the WUI; two classes of WUI were identified:

- the intermix, where there is at least 50% vegetation cover surrounding buildings
- the interface, where buildings are within 2.4 km (1.5 miles) of a patch of vegetation at least 5 km<sup>2</sup> in size that contains at least 75% vegetation.

Both classes required a minimum building density of 6.17 buildings per km<sup>2</sup> (using a range of circular neighborhood sizes).

This is a proprietary index developed by Pyrologix, representing the level of wildfire exposure for a structure (e.g., a home) if one were to exist on a given pixel. It is an integrated measure that includes three components: the likelihood of a wildfire of any intensity occurring in a given year (annual burn probability), potential wildfire intensity for a given pixel, and ember load to that pixel from surrounding vegetation.

SES data was produced by Pyrologix LLC, a wildfire threat assessment research firm, as part of a spatial wildfire hazard assessment across all land ownerships for the state of California. The ongoing work generally follows the framework outlined in Scott and Thompson (2013), with custom methods and significant improvements developed by Pyrologix. The project generally consists of three components: fuelscape calibration and updates, wildfire hazard assessment, and risk assessment. It utilizes a combination of wildfire models and custom tools, including the FSim large wildfire simulator (Finney et al., 2011), and WildEST, a custom modeling tool developed by Pyrologix (Scott, 2020). To date, this work has resulted in a wide variety of spatial data layers related to wildfire hazard and risk, including Structure Exposure Score (SES), representing conditions prior to the 2020, 2021 and 2022 fire seasons. Work to date has been funded by the USDA Forest Service Region 5, the California Energy Commission, and the USDI Bureau of Land Management with data contributions from CAL FIRE.

For this project, the FSim large-fire simulator is used to quantify annual wildfire likelihood across the analysis area. FSim is a comprehensive fire occurrence, growth, behavior, and suppression simulation system that uses locally relevant fuel, weather, topography, and historical fire occurrence information to make a spatially resolved estimate of the contemporary likelihood and intensity of wildfire across the landscape.

WildEST (Wildfire Exposure Simulation Tool) is used to quantify wildfire intensity and ember loads across the analysis area. WildEST is a deterministic wildfire modeling tool developed by Pyrologix that integrates spatially continuous weather input variables, weighted based on how they will likely be realized on the landscape. This

makes the deterministic intensity values developed with WildEST more robust for use in effects analysis than the stochastic intensity values developed with FSim. This is especially true in low wildfire occurrence areas where predicted intensity values from FSim are reliant on a very small sample size of potential weather variables. It also allows for more appropriate weighting of high-spread conditions into fire behavior calculations. WildEST also produces indices of conditional and expected ember production from vegetated areas (pixels) and load to other pixels in the analysis area. Please reference the Pyrologix 2021 project report (Volger et al., 2021) for more information on WildEST analysis.

FSim was run for the CAL 2022 fuelscape at 120m resolution. WildEST was run for the CAL 2022 fuelscape at 30-m resolution. Both models utilized gridded hourly historical California weather data provided by CALFIRE. Results for annual burn probability (FSim), fire intensity (WildEST) and ember load (WildEST) were used to create Structure Exposure Score.

**Data Source:**

Pyrologix, LLC

WUI, Carlson et al, 2022

**File Name:** StructureExposureScore\_WUI\_2022.tif

---

## DAMAGE POTENTIAL

**Tier:** 1

**Data Vintage:** 2022

**Metric Definition and Relevance:** This metric combines two data layers; one is the Wildland Urban Interface (WUI) as defined by Carlson et al. 2022 (see [WUI definition](#) section for more information), and a second data layer, Damage Potential (DP), developed by Pyrologix LLC. The WUI includes the intermix and interface zones which collectively identify areas where structures occur. The distance selected for the interface definition is based on research from the California Fire Alliance suggesting that this is the average distance firebrands can travel from an active wildfire front.

The composite Damage Potential (DP) dataset represents a relative measure of wildfire’s potential to damage a home or other structure if one were present at a given pixel, and if a wildfire were to occur (conditional exposure). It is a function of ember load to a given pixel, and fire intensity at that pixel, and considers the generalized consequences to a home from fires of a given intensity (flame length). This index does not incorporate a measure of annual wildfire likelihood.

**Data Resolution:** 30m Raster

**Data Units:** Relative index, low to high

**Creation Method:** This metric represents DP for WUI areas only. DP values were binned based on the following ranges into 6 classes and assigned class names.

- 0 (None): Values = 0
- 1 (Very Low): Values 0.01 to 20
- 2 (Low): Values 20 to 35
- 3 (Moderate): Values 35 to 50
- 4 (High): Values 50 to 80
- 5 (Very High): Values 80+

The current delineation of the WUI (Carlson et al. 2022) uses a mapping algorithm with definitions of the WUI; two classes of WUI were identified:

- the intermix, where there is at least 50% vegetation cover surrounding buildings
- the interface, where buildings are within 2.4 km (1.5 miles) of a patch of vegetation at least 5 km<sup>2</sup> in size that contains at least 75% vegetation.

Both classes required a minimum building density of 6.17 buildings per km<sup>2</sup> (using a range of circular neighborhood sizes).

Damage Potential (DP) data was produced by Pyrologix LLC, a wildfire threat assessment research firm, as part of a spatial wildfire hazard assessment across all land ownerships for the state of California. The ongoing work generally follows the framework outlined in Scott and Thompson (2013), with custom methods and significant improvements developed by Pyrologix. The project generally consists of three components: fuelscape calibration and updates, wildfire hazard assessment, and risk assessment. It utilizes a combination of wildfire models and custom tools, including WildEST (Wildfire Exposure Simulation Tool), a custom modeling tool developed by Pyrologix (Scott, 2020). To date, this work has resulted in a wide variety of spatial data layers related to wildfire hazard and risk, including Damage Potential (DP), representing conditions prior to the 2020, 2021 and 2022 fire seasons. Work to date has been funded by the USDA Forest Service Region 5, the California Energy Commission, and the USDI Bureau of Land Management with data contributions from CAL FIRE. Please reference the Pyrologix 2021 project report (Volger et al., 2021) for more information about the project or WildEST analysis.

Damage Potential (DP) is a proprietary index developed by Pyrologix LLC representing wildfire's potential to damage a home or other structure if a wildfire were to occur (conditional exposure). It is a function of ember load to a given pixel and fire intensity at that pixel, and it considers the generalized consequences to a home from fires of a given intensity (flame length). DP is calculated based on two other datasets developed by Pyrologix: conditional risk to potential structures (cRPS) and conditional ember load index (cELI).

cRPS represents the potential consequences of fire to a home at a given location if a fire occurs there and if a home were located there. It is a measure that integrates wildfire intensity with generalized consequences to a home on every pixel. Wildfire intensity (flame length) is calculated using Pyrologix' WildEST tool. WildEST is a scripted geospatial process used to perform multiple deterministic simulations under a range of weather types (wind speed, wind direction, fuel moisture content). Rather than weighting results solely according to the temporal relative frequencies of the weather scenarios, the WildEST process integrates results by weighting them according to their weather type probabilities (WTP), which appropriately weights high-spread conditions into the calculations. For fire-effects calculations, WildEST generates flame-length probability rasters that incorporate non-heading spread directions, for which fire intensity is considerably lower than at the head of the fire.

The response function characterizing potential consequences to an exposed structure is applied to fire effects flame lengths from WildEST for all burnable fuel types on the landscape regardless of whether an actual structure is present or not. The response function does not consider building materials of structures and is meant as a measure of the effect of fire intensity on structure exposure. The response function is provided below:

- Flame length probability of 0-2 ft: -25
- Flame length probability of 2-4 ft: -40
- Flame length probability of 4-6 ft: -55
- Flame length probability of 6-8 ft: -70
- Flame length probability of 8-12 ft: -85
- Flame length probability of >12 ft: -100

These results were calculated using 30m fire-effects flame-length probabilities from the WildEST wildfire behavior results and then further smoothed.

cELI is also calculated in WildEST, and represents the relative ember load per pixel, given that a fire occurs, based on surface and canopy fuel characteristics, climate, and topography within the pixel. Units are the relative number of embers. cELI is based on heading-only fire behavior.

Damage Potential is then calculated as the arithmetic mean of cELI and cRPS for each pixel across the landscape as follows:

$$DP = cRPS + cELI/2$$

Although flame length and its potential impact to structures is a function of the fire environment at the subject location only, ember load is a function of ember production and transport in the area surrounding the subject location. A location with light fuel (and therefore low flame length) could still have significant Damage Potential if surrounded by a fire environment that produces copious embers.

**Data Source:**

Pyrologix, LLC

WUI, Carlson et al, 2022

**File Name:** DamagePotential\_WUI\_2022.tif

---

EMBER LOAD INDEX

**Tier:** 1

**Data Vintage:** 2022

**Metric Definition and Relevance:** This ember load dataset represents the ember load index (ELI) per pixel, for a given pixel, based on surface and canopy fuel characteristics, climate, and topography within the pixel. The Ember Load Index (ELI) incorporates burn probability (BP). BP is incorporated into calculations of the ember production before the distribution of embers across the landscape to determine ember load. Given that ELI incorporates burn probability, this index can be used to identify where on the landscape hardening buildings may be needed to resist ignition and the priority for doing so according to the likelihood of the area being visited by fire.

**Data Resolution:** 30m Raster

**Data Units:** Relative number of embers.

**Creation Method:** ELI is not simply the multiplication of ember load (ELI) and burn probability (BP). Rather, BP is incorporated into calculations of the ember production prior to the distribution of embers across the landscape to determine ember load. ELI is based on heading-only fire behavior.

**Data Source:** Pyrologix, LLC

**File Name:** EmberLoadIndex\_2022.tif

---

IGNITION CAUSE

**Tier:** 1

**Data Vintage:** 1992 - 2020

**Metric Definition and Relevance:** The original point layer (WildfireOccurrence\_CA\_1992\_2020.shp ) contains a spatial database of wildfires that occurred in the United States from 1992 to 2020. It is the fifth update of a publication originally generated to support the national Fire Program Analysis (FPA) system. The wildfire records were acquired from the reporting systems of federal, state, and local fire organizations. The following core data elements were required for records to be included in this data publication: discovery date, final fire size, and a point location at least as precise as a Public Land Survey System (PLSS) section (1-square mile grid). The data were transformed to conform, when possible, to the data standards of the National Wildfire Coordinating Group (NWCG), including an updated wildfire-cause standard (approved August 2020). Basic error-checking was performed and redundant records were identified and removed, to the degree possible. The resulting product, referred to as the Fire Program Analysis fire-occurrence database (FPA FOD), includes 2.3 million geo-referenced wildfire records, representing a total of 180 million acres burned during the 29-year period. Identifiers necessary to link the point-based, final-fire-reporting information to published large-fire-perimeter and operational-situation-reporting datasets are included. Short, Karen C. 2022. Spatial wildfire occurrence data for the United States, 1992-2020 [FPA\_FOD\_20221014]. 6th Edition. Fort Collins, CO: Forest Service Research Data Archive. <https://doi.org/10.2737/RDS-2013-0009.6>

**Data Resolution:** Vector(points) and 30m Raster

**Data Units:** Categorical

**Creation Method:** Rocky Mountain Research Station (U.S. Forest Service) scientist, Karen Short, is the principal creator of this data set. Points were converted to 30m raster cells using the “most frequent” function on the NWCG\_CAUSE\_CLASSIFICATION attribute (Broad classification of the reason the fire occurred) creating three rasters:

- Human caused ignition
- Lightning (natural) caused ignition
- All causes of ignition - Human or Natural and Missing data/not specified/undetermined

“MostFrqCau” indicates the most frequent cause of the fire in that location.. “FireCount” indicates the number of fires that occurred between 1992 and 2020, regardless of cause. It is noted that locations with hundreds of counts may be a result of the method of how ignitions are reported/recorded. Both the accuracy and precision of the location estimates are generally much lower than that implied by the stored coordinate information – which, for example, may have been calculated from a PLSS section centroid. Efforts were made to purge redundant records to the best of the authors’ ability. Despite this, some locations may have multiple records that may reflect redundant records or multiple reports of fires due to the imprecision of the location record, the reporting process of an individual authority, or the possible reality of multiple initiations at a given location.

**Data Source:** Rocky Mountain Research Station, U.S. Forest Service

File Name: WldfireAllCausesCount\_1992\_2020.tif; WldFireOccCause\_Human\_1992\_2020.tif;  
WldFireOccCause\_Natural\_1992\_2020.tif; WildfireOccurrence\_CA\_1992\_2020.shp

---

## SOURCE OF EMBER LOAD TO BUILDINGS

**Tier:** 1

**Data Vintage:** 2022

**Metric Definition and Relevance:** The ember transport model used in WildEST tracks the travel of embers from each source pixel to downwind receiving pixels. The relative number of embers landing on a given receiving pixel is summed across all potential source pixels. If the receiving pixel has a nonzero WRC Building Cover value (meaning the pixel is within 75 m of a qualifying building), then we separately sum the relative number of embers from the source pixel. The final SELB raster represents the expected annual relative ember production that lands on building cover across all weather types.

**Data Resolution:** 30m Raster

**Data Units:** Relative index

**Creation Method:** The WildEST modeling contains a module for producing indices of conditional and expected ember production and load. The Conditional Ember Production Index (cEPI) is an index of the relative number of embers lofted at a given landscape pixel given the fire environment there, given that a fire occurs. Ember Production Index (EPI) is the expected value of cEPI; it is the expected annual relative number of embers lofted from a given landscape pixel.

The Conditional Ember Load Index (cELI) is a relative index of the relative number of embers that land at a given landscape location, including nonburnable pixels. Finally, Ember Load Index combines the conditional ELI and the likelihood of that ember load occurring. All ember characteristics are based on headfire behavior. These

The ember load indices represent relative ember load at a pixel. Similar to ember production, ember load is also based on surface and canopy fuel characteristics, climate, and topography at the pixel. Ember load incorporates downwind ember travel.

The Ember Load Index (ELI) incorporates burn probability; however, ELI is not simply the multiplication of condition ember load (cELI) and burn probability (BP). Rather, BP is incorporated into calculations of the ember production before the distribution of embers across the landscape to determine ember load. Given that ELI incorporates burn probability, this index can be used to identify where on the landscape hardening buildings may be needed to resist ignition and the priority for doing so according to the likelihood of the area being visited by fire.

**Data Source:** Pyrologix, LLC

**File Name:** SourceEmberLoadToBuildings.tif

---

## WILDFIRE HAZARD POTENTIAL

**Tier:** 1

**Data Vintage:** 2022

**Metric Definition and Relevance:** Wildfire Hazard Potential (WHP) is an index that quantifies the relative potential for wildfire that may be difficult to control. WHP can be used as a measure to help prioritize where fuel treatments may be needed.

**Data Resolution:** 30m Raster

**Data Units:** Relative index

**Creation Method:** Pyrologix calculated WHP following the methods established by Dillon et al. (2015) and Dillon (2018). The original methods utilize lower-resolution FSim inputs, while our approach uses higher-resolution inputs



including 30-m CAL vegetation inputs (derived from LANDFIRE 2016), 30-m CAL fuel model outputs, 30-m CAL burn probability results, and 30-m CAL fire-effects flame-length probabilities from the WildEST wildfire behavior results.

**Data Source:** Pyrologix, LLC

**File Name:** WildfireHazardPotential\_2022.tif

## FIRE DYNAMICS

Fire dynamics reflect fire as an ecological process and the function that it performs. It can be broken into two key elements: functional fire and fire severity. Although fire dynamics pertain to the entire landscape, the ecological role of fire is most relevant to landscapes outside of the wildland urban interface (WUI). Within the WUI, protection of life and property takes priority over the role of fire as a process. As a result, this fire dynamics pillar pertains largely to areas outside of the WUI while the fire-adapted communities pillar pertains largely to areas inside the WUI.

**DESIRED OUTCOME:** Fire burns in an ecologically beneficial and socially acceptable way that perpetuates landscape heterogeneity and rarely threatens human safety or infrastructure.

## FUNCTIONAL FIRE

Increasing the pace and scale of restoration on the landscape will require using a variety of tools to accomplish restoration targets. The use of prescribed fire and managed wildfires, where appropriate, can contribute to the restoration need. This is particularly true where fires burn at low and moderate severity, which we are referring to as “functional fire”. Functional fire is when fire burns in an ecologically beneficial and socially acceptable way, perpetuating landscape heterogeneity and rarely threatening human safety or infrastructure.

### **Updated FVEG; Methods for 2023 statewide updates to FVEG WHRtype, WHRsize, and WHRdensity**

An accurate depiction of the spatial distribution of vegetation/habitat types within California is required for a number of the metrics included in this kit, particularly for some of the fire, forest and rangeland resiliency, and biodiversity metrics. The California Department of Forestry and Fire Protections CALFIRE Fire and Resource Assessment Program (FRAP), in cooperation with California Department of Fish and Wildlife VegCamp program and extensive use of USDA Forest Service Region 5 Mapping and Remote Sensing unit (MARS) data, has compiled the “best available” land cover data available for California into a single comprehensive statewide data set. The data span a period from approximately 1990 to 2014.

Because the source data are in many cases fairly old and there has been extensive disturbance, particularly from wildfire, over the last 25 years, we made some updates to the 2015 version of FVEG. The methods for making those changes are described here.

#### *WHRtype update*

FVEG’s WHRtype was updated with the LANDFIRE Existing Vegetation Type (EVT) data product version 2.2.0 (LANDFIRE 2020) and the Rangeland Analysis Platform (RAP) fractional ground cover data product version 3.0 (Jones et al. 2018, Allred et al. 2021). Pixels were considered for update where high severity wildfire occurred after the FVEG mapping date. High severity was defined as wildfire burned areas that experienced  $\geq 75\%$  loss in basal

area (Parks et al. 2018, Young-Hart et al. 2022) following the wildfire event. The type of update that occurred in each “high severity” pixel was dependent upon a lifeform conversion comparison (FVEG-to-LANDFIRE EVT), vegetation height (SALO 2020), and percent ground cover by annual and perennial grasses (RAP) (Table 1).

Table 1. FVEG-LANDFIRE update type for high severity pixels. Annual grass (AG) cover and perennial grass (PG) cover data were from the Rangeland Analysis Platform fractional ground cover data product version 3.0. Canopy height (CH) data were from the SALO forest observatory data product.

|                                      |                   | FVEG Lifeform  |  |  |
|--------------------------------------|-------------------|--|--|--|
|                                      |                   | <i>Herbaceous</i>  | <i>Shrub</i>   | <i>Tree</i>  |
| <b>LANDFIRE<br/>EVT<br/>Lifeform</b> | <i>Herbaceous</i> | <ul style="list-style-type: none"> <li>Where AG cover &gt;50%, update to “Annual Grassland” <u>WHRname (“AGS” WHRtype)</u></li> </ul>    | <ul style="list-style-type: none"> <li>Where CH &lt;1m &amp; AG cover &gt;50%, update to “Annual Grassland” <u>WHRname (“AGS” WHRtype)</u></li> </ul>    | <ul style="list-style-type: none"> <li>Where CH &lt;2m &amp; AG cover &gt;50%, update to “Annual Grassland” <u>WHRname (“AGS” WHRtype)</u></li> </ul>    |
|                                      |                   | <ul style="list-style-type: none"> <li>Where PG cover &gt;50%, update to “Perennial Grassland” <u>WHRname (“PGS” WHRtype)</u></li> </ul> | <ul style="list-style-type: none"> <li>Where CH &lt;1m &amp; PG cover &gt;50%, update to “Perennial Grassland” <u>WHRname (“PGS” WHRtype)</u></li> </ul> | <ul style="list-style-type: none"> <li>Where CH &lt;2m &amp; PG cover &gt;50%, update to “Perennial Grassland” <u>WHRname (“PGS” WHRtype)</u></li> </ul> |
|                                      |                   | <ul style="list-style-type: none"> <li>Otherwise, no update to <u>WHRtype</u></li> </ul>   | <ul style="list-style-type: none"> <li>Otherwise, FVEG-LANDFIRE crosswalk update</li> </ul>  | <ul style="list-style-type: none"> <li>Otherwise, FVEG-LANDFIRE crosswalk update</li> </ul>  |
|                                      | <i>Shrub</i>      | FVEG-LANDFIRE crosswalk update   | No update to <u>WHRtype</u>  | FVEG-LANDFIRE crosswalk update   |
|                                      | <i>Tree</i>       | FVEG-LANDFIRE crosswalk update   | <ul style="list-style-type: none"> <li>FVEG-LANDFIRE crosswalk update</li> </ul>   | No update to <u>WHRtype</u>  |

*WHRdensity and WHRsize updates*

Following the WHRtype update, pixels that had lifeform “tree” then had the FVEG attributes “WHRdensity” and “WHRsize” updated using the SALO Forest Observatory canopy height and canopy cover data products (SALO 2020). SALO data were available for the years 2016-2020, values of canopy height and canopy cover were averaged across years for the update[2].

To update WHRdensity, SALO canopy cover was converted to WHRdensity canopy closure class per the [Wildlife Habitat Relationships, Standards for Canopy Closure Table 114C](#).

To update WHRsize, we developed allometric equations that predict tree DBH (diameter at breast height, breast height = 4.5 ft) as a function of tree height (HT, ft). We used data from the USDA Forest Inventory and Analysis program (FIA) for California (FIA DataMart 2023; California 2022 database; ver. 9.0.1). For this analysis, we included live trees ≥ 4.5 ft tall with a crown class code of dominant, co-dominant, or open grown (N = 165,224 tree measurements between 1991 and 2019). Trees were grouped by region based on the “fuzzed” location of the plot. Regions were defined by the Regional Resource Kits (2023, 4 regions) and separated into softwoods and hardwoods as defined by FIA (2 categories). For each analysis, three functions were evaluated: linear, saturating, and power:

Linear:  $DBH = a + (b*HT)$ ;

Saturating (Michaelis–Menten):  $DBH = (V_m * HT) / (K + HT)$ ;

Power:  $DBH = aHT^b$ .

For the most informative model (i.e., lowest AIC), we report both the root mean squared error (RMSE) and the pseudo  $R^2$ . In this case, pseudo  $R^2$  was calculated as the coefficient of determination between the observed and predicted DBHs (Table 2). We used the most informative HT-to- DBH function for the region and tree category to convert SALO canopy height data to DBH that was then converted to WHRsize class per the [Wildlife Habitat Relationships, Standards for Tree Size Table 114B](#).

Table 2. Height-to-DBH conversion equations by California region and tree class. DBH is in inches; Height (HT) is in feet. Only included trees with a HT > = 4.5 feet. Only included canopy class = dominant, co-dominant, or open. Equation (EQN) code: MM = Michaelis Menton; POWER = power; Linear = linear.

| Region              | Tree Class | EQN    | a (Vm) | b (K)  | RMSE | pseudoR <sup>2</sup> | EQN formula                   |
|---------------------|------------|--------|--------|--------|------|----------------------|-------------------------------|
| Sierra Nevada       | Softwood   | MM     | 223.39 | 712.20 | 6.57 | 0.69                 | $DBH = (V_m * HT) / (K + HT)$ |
| Sierra Nevada       | Hardwood   | Linear | -0.391 | 0.294  | 4.69 | 0.57                 | $DBH = a + b(HT)$             |
| Southern California | Softwood   | MM     | 108.97 | 216.30 | 7.47 | 0.55                 | $DBH = (V_m * HT) / (K + HT)$ |
| Southern California | Hardwood   | MM     | 175.17 | 424.31 | 5.55 | 0.52                 | $DBH = (V_m * HT) / (K + HT)$ |
| North Coast         | Softwood   | POWER  | 0.128  | 1.13   | 6.51 | 0.74                 | $DBH = a * HT^b$              |
| North Coast         | Hardwood   | Linear | 0.135  | 0.242  | 5.2  | 0.49                 | $DBH = a + b(HT)$             |
| Central Coast       | Softwood   | Linear | 0.588  | 0.244  | 8.25 | 0.62                 | $DBH = a + b(HT)$             |
| Central Coast       | Hardwood   | MM     | 68.51  | 161.40 | 6.24 | 0.45                 | $DBH = (V_m * HT) / (K + HT)$ |

### Availability of Data and Materials

Data used for the 2023 FVEG updates can be obtained from the following:

- LANDFIRE – <http://www.landfire.gov/>
- Rangeland Analysis Platform – <https://rangelands.app/products/>
- SALO Forest Canopy – <https://forestobservatory.com/download>
- 10-year summary of basal area lost – <https://data.fs.usda.gov/geodata/rastergateway/acre>
- Perturbed FIA data – <https://experience.arcgis.com/experience/3641cea45d614ab88791aef54f3a1849>

Google Earth Engine Python API script can be obtained from: [https://github.com/kjohnston73/fveg\\_update](https://github.com/kjohnston73/fveg_update)

### References

Allred, B.W., B.T. Bestelmeyer, C.S. Boyd, C. Brown, K.W. Davies, M.C. Duniway, L.M. Ellsworth, T.A. Erickson, S.D. Fuhlendorf, T.V. Griffiths, and V. Jansen. (2021). Improving Landsat predictions of rangeland fractional cover with multitask learning and uncertainty. *Methods in Ecology and Evolution* 12 (5): 841–849.

Burnham, K.P., and D.R. Anderson. 2002. *Model selection and multimodel inference: a practical information-theoretic approach*. 2nd ed. New York, Springer-Verlag.

FIA DataMart. 2023. USDA Forest Inventory and Analysis DataMart. <https://experience.arcgis.com/experience/3641cea45d614ab88791aef54f3a1849/>

Jones, M.O., B.W. Allred, D.E. Naugle, J.D. Maestas, P. Donnelly, L.J. Metz, J. Karl, R. Smith, B. Bestelmeyer, C. Boyd, and J.D. Kerby. (2018). Innovation in rangeland monitoring: annual, 30 m, plant functional type percent cover maps for US rangelands, 1984–2017. *Ecosphere* 9 (9): 02430.

LANDFIRE: LANDFIRE Existing Vegetation Type layer. (2020). U.S. Department of Interior, Geological Survey, and U.S. Department of Agriculture.

Parks, S.A., Holsinger, L.M., Voss, M.A., Loehman, R.A., Robinson, N.P. (2018). Mean composite fire severity metrics computed with Google Earth Engine offer improved accuracy and expanding mapping potential. *Remote Sensing*. v10(6), 879.

Regional Resource Kits. 2023. California Wildfire and Forest Resilience Task Force. <https://wildfiretaskforce.org/regional-resource-kits-page/>

SALO. (2020). <https://salo.ai/assets/pdf/Forest-Observatory-Data-Description.pdf>.

Young-Hart L., Stine P., Manley P., Clark C., Bistriz L., Goulden M., Kuskulis E., Ramirez C., Slaton M., Stewart J.A.E, Tarnay L., Young, D.J. (2022) ACCEL Metric Dictionary Version 3.0. USDA Forest Service.

---

### FIRE IGNITION PROBABILITY

**Tier: 3**

**Data Vintage:** 1992 to 2015

**Metric Definition and Relevance:** These rasters depict the predicted human- and lightning-caused ignition probability for the state of California.

**Data Resolution:** 1km Raster

**Data Units:** Probability, 0-1

**Creation Method:** [Spatial patterns and drivers for wildfire ignitions in California - IOPscience](#)

**Data Source:** Bin Chen and Yufang Jin, University of California Davis, bch@ucdavis.edu

**File Name:** PredictedHumanIgnitionProb\_1km.tif; PredictedLightningIgnitionProb\_1km.tif

## FIRE RETURN INTERVAL DEPARTURE

**Definition and Relevance:** The fire return interval departure (FRID) analysis quantifies the difference between current and pre-settlement fire frequencies, allowing managers to target areas at high risk of threshold-type responses owing to altered fire regimes and interactions with other factors.

**Creation Method:** The FRID methodology was developed and described by Van de Water and Safford (2011). The feature class is now produced and maintained by the U.S. Forest Service, Region 5, Information Management – Mapping and Remote Sensing (MARS) Team. Contemporary FRIs were calculated using the fire dates and footprints from California Interagency Fire Perimeters database (maintained by the California Department of Forestry and Fire Protection (CalFire-FRAP). The vegetation type stratification (i.e. to calculate the FRI for individual vegetation types) was based on the MARS Existing Vegetation (EVEG) map for California from the year 2011, with the vegetation typing (“CALVEG”) cross-walked (grouped) into 28 pre-settlement fire regime (PFR) types.

For assorted reasons, portions of San Benito and San Luis Obispo Counties never received a full EVEG Baseline Mapping assessment and thus data in the FRID Central Coast layer has some holes in these areas. In 2009, an EVEG mapping project was started for these areas but never finalized. San Luis Obispo County, the southern part of Santa Clara County, and all of San Benito County were baseline mapped using the Hardwood Dataset as a foundation for regional dominance (vegtype). Additional data sources from the National Land Cover Database, San Luis Obispo County Farm Data, Farmland Mapping & Monitoring Program, Bureau of Reclamation, and National Hydrology Database were then used to overwrite the Hardwood data where it was relevant. Structural attributes for forested conditions came primarily from the Hardwoods Dataset for canopy values while tree size was derived from a classification of Thematic Mapper 30-meter imagery.

Although incomplete as an EVEG database, these “best available data” were used by the RRK team to fill holes in FRID for the Central Coast RRK project. The MARS team completed a crosswalk from Regional Dominance Type 1 (vegtype) to the FRID PFR attribute and calculations for the “gap” areas were run for fire return interval departure. We have used this “patch” to address FRID needs for the near-term. The data for these areas will show vulnerabilities to analysis at larger scales until a time that these areas can be visually edited to match the level of precision seen in the adjoining Los Padres NF.

Other gaps (NoData):

Although areas mapped as grasslands and meadows were included in the GIS layer, FRI and departure statistics were not calculated for these types because reliable information about pre-Euroamerican settlement fire regimes is lacking. These values (-999) have been converted to NoData in the RRK datasets.

**Data Source:** USDA Forest Service, Region 5, MARS Team

**References:** Information on pre-Euromerican settlement FRIs (fire return intervals) was compiled from an exhaustive review of the fire history literature, expert opinion, and vegetation modeling (Van de Water and Safford 2011; Safford and Van de Water 2014). Contemporary FRIs were calculated using the California Interagency Fire Perimeters database (maintained by the California Department of Forestry and Fire Protection (CAL FIRE-FRAP). The vegetation type stratification was based on the US Forest Service existing vegetation map (USDA Forest Service, Mapping and Remote Sensing Team) for California from the year 2011, with the vegetation typing (“CALVEG”) grouped into 28 pre-settlement fire regime (PFR) types, as defined by Van de Water and Safford (2011). The 2011 eVeg map is used as the baseline for all subsequent FRID maps to freeze the underlying vegetation template and permit temporal comparisons without introducing vegetation type change as a confounding factor.

---

## MEAN PERCENT FRI DEPARTURE, SINCE 1908

**Tier:** 3

**Data Vintage:** 2021

**Metric Definition and Relevance:** This metric, mean percent FRID, is a measure of the extent to which contemporary fires (i.e., since 1908) are burning at frequencies similar to the frequencies that occurred prior to Euro-American settlement, with the mean reference FRI as the basis for comparison. Mean PFRID is a metric of fire return interval departure (FRID) and measures the departure of current FRI from mean reference FRI in percent.

**Data Resolution:** 30m Raster

**Data Units:** Percent

**Creation Method:** The current FRI is calculated by dividing the number of years in the fire record (e.g., 2019-1908=112 years inclusive) by the number of fires occurring between 1908 and the current year in a given polygon plus one ( $\text{CurrentFRI} = \text{Number of years} / \text{Number of fires} + 1$ ). The mean reference FRI is an approximation of how often, on average, a given PFR likely burned in the three or four centuries prior to significant Euro-American settlement. This measure does not return to zero when a fire occurs, unlike FRID values used in some other analyses (e.g., NPS FRID Index). Instead, the following formulas are used to calculate Mean PFRID:

When current FRI is longer than reference FRI (the common condition in most coniferous PFRs) the formula is:

$$[1 - (\text{MeanRefFRI} / \text{CurrentFRI})] * 100$$

When current FRI is shorter than reference FRI (common in some shrub dominated PFRs, and areas in the Wildland Urban Interface) the formula is:

$$- \{ [1 - (\text{CurrentFRI} / \text{MeanRefFRI})] \} * 100$$

For areas dominated by PFRs with a mean reference FRI greater than 112 years, and that have not burned in the period of historical record considered in this analysis (i.e., since 1908), the FRID is assumed to equal zero.

**Data Source:**

Fire History (2021), CAL FIRE

Existing Vegetation (CALVEG 2011), USDA Forest Service, Region 5, MARS Team

**File Name:** CenCst\_meanPFRID.tif

---

## MEAN PERCENT FRI DEPARTURE, SINCE 1970

**Tier: 2**

**Data Vintage: 2021**

**Metric Definition and Relevance:** Mean Percent FRID (meanPFRID\_1970) quantifies the extent in percentage to which recent fires (i.e., since 1970) are burning at frequencies similar to those that occurred prior to Euro-American settlement, with the mean reference FRI as the basis for comparison. Mean PFRID measures the departure of current FRI from reference mean FRI in percent.

**Data Resolution: 30m Raster**

**Data Units: Percent**

**Creation Method:** The current FRI is calculated by dividing the number of years in the fire record (e.g., 2019-1970=49 years inclusive) by the number of fires occurring between 1970 and the current year in a given polygon plus one ( $\text{CurrentFRI} = \text{Number of years} / \text{Number of fires} + 1$ ). The mean reference FRI is an approximation of how often, on average, a given PFR likely burned in the three or four centuries prior to significant Euro-American settlement. This measure does not return to zero when a fire occurs, unlike FRID values used in some other analyses (e.g., NPS FRID Index). The following formulas are used to calculate Mean PFRID, the same as with meanPFRID but with 1970 as the baseline rather than 1908. Important note: because 1970 is the baseline for this measure, no fires before 1970 are taken into account and all PFRs start at a PFRID of zero beginning in 1970.

When current FRI is longer than reference FRI (the common condition in most coniferous PFRs) the formula is:

$$[1 - (\text{MeanRefFRI} / \text{CurrentFRI})] * 100$$

When current FRI is shorter than reference FRI (common in some shrub dominated PFRs, and areas in the Wildland Urban Interface) the formula is:

$$- \{ [1 - (\text{CurrentFRI} / \text{MeanRefFRI})] \} * 100$$

**Data Source:**

Fire History (2021), CAL FIRE

Existing Vegetation (CALVEG 2011), USDA Forest Service, Region 5, MARS Team

**File Name:** CenCst\_meanPFRID\_1970.tif

---

## FRID CONDITION CLASS FOR DEPARTURE

**Tier: 3**

**Data Vintage: 2021**

**Metric Definition and Relevance:** This metric uses the mean percent FRID to a measure of the extent to which contemporary fires (i.e., since 1908) are burning at frequencies similar to the frequencies that occurred prior to Euro-American settlement, with the mean reference FRI binned into another basis for comparison. Mean PFRID is a

metric of fire return interval departure (FRID), and measures the departure of current FRI from reference mean FRI in percent.

**Data Resolution:** 30m Raster

**Data Units:** Integer, -3 to 3

**Creation Method:** This is a condition class categorization of the data in the Mean PFRID field. MeanCC\_FRI categorizes the percent differences calculated in Mean PFRID using the following scale:

- 1      0 to 33.3%
- 2      33 to 66.7%
- 3      greater than 66.7%

Negative condition classes (i.e., where fires are burning more often than under pre-Anglo-American settlement conditions) are categorized on the negative of the same scale:

- 1      0 to -33.3%
- 2      -33 to -66.7%
- 3      less than -66.7%

CC1 and CC-1 are mapped in the same class because they are both within 33% of the mean pre-settlement value.

**Data Source:**

Fire History (2021), CAL FIRE

Existing Vegetation (CALVEG 2011), USDA Forest Service, Region 5, MARS Team

**File Name:** CenCst\_meanCC\_FRI.tif

---

## TIME SINCE LAST FIRE

**Tier:** 2

**Data Vintage:** 2021

**Metric Definition and Relevance:** Time Since Last Fire (TSLF), from the Fire Return Interval Departure (FRID) map, provides information (in years) to indicate the length of time since an area last burned.

**Data Resolution:** 30m Raster

**Data Units:** Years

**Creation Method:** Time Since Last Fire (TSLF), from the Fire Return Interval Departure (FRID) map, provides information (in years) to indicate the length of time since an area last burned. Specifically, the number of years elapsed between the most recent fire recorded in the fire perimeters database and the version year of the FRID map being used. To illustrate, if the version year of the FRID map is 2019, and the area in question last burned in 1995, TSLF will be 24 (2019 minus 1995).



**Data Source:**

Fire History (2021), CAL FIRE  
Existing Vegetation (CALVEG 2011), USDA Forest Service, Region 5, MARS Team

**File Name:** CenCst\_TSLF.tif

---

**CURRENT FIRE RETURN INTERVAL DEPARTURE, SINCE 1908**

**Tier:** 3

**Data Vintage:** 2021

**Metric Definition and Relevance:** The fire return interval departure (FRID) analysis quantifies the difference between current and pre-settlement fire frequencies, allowing managers to target areas at high risk of threshold-type responses owing to altered fire regimes and interactions with other factors. This is a measure of the extent to which contemporary fires (i.e. since 1908) are burning at frequencies similar to the frequencies that occurred prior to Euro-American settlement.

**Data Resolution:** 30m Raster

**Data Units:** Average Years

**Creation Method:** Current fire return interval 1908 is calculated by dividing the number of years in the fire record by the number of fires occurring between 1908 and the current year in a given polygon plus one.

$$\text{CurrentFRI} = \text{Number of years} / \text{Number of Fires} + 1$$

**Data Source:**

Fire History (2021), CAL FIRE  
Existing Vegetation (CALVEG 2011), USDA Forest Service, Region 5, MARS Team

**File Name:** CenCst\_currentFRI.tif

---

**CURRENT FIRE RETURN INTERVAL DEPARTURE, SINCE 1970**

**Tier:** 3

**Data Vintage:** 2021

**Metric Definition and Relevance:** The fire return interval departure (FRID) analysis quantifies the difference between current and pre-settlement fire frequencies, allowing managers to target areas at high risk of threshold-type responses owing to altered fire regimes and interactions with other factors. This is a measure of the extent to which contemporary fires (i.e. since 1970) are burning at frequencies similar to the frequencies that occurred prior to Euro-American settlement, with the mean reference FRI as the basis for comparison. With this metric, mPFRID\_1970, the same formulas are used as with meanPFRID but with 1970 as the baseline rather than 1908. Important note: because 1970 is the baseline for this measure, no fires before 1970 are taken into account and all PFRs start at a PFRID of zero beginning in 1970.

**Data Resolution:** 30m Raster

**Data Units:** Average Years

**Creation Method:** Current fire return interval 1970 is calculated by dividing the number of years in the fire record by the number of fires occurring between 1970 and the current year in a given area plus one.

$$\text{CurrentFRI}_{1970} = \text{Number of years} / \text{Number of Fires} + 1$$

**Data Source:**

Fire History (2021), CAL FIRE  
Existing Vegetation (CALVEG 2011), USDA Forest Service, Region 5, MARS Team

**File Name:** CenCst\_currentFRI\_1970.tif

**SEVERITY**

Uncharacteristic proportions of high severity fire over the area burned, particularly in the last decade, has been a common theme in the megafires that have occurred throughout the Central Coast Region recently. The following metrics characterize, map, and quantify some of the factors that contribute.

---

**ANNUAL BURN PROBABILITY**

**Tier:** 1

**Data Vintage:** 2022

**Metric Definition and Relevance:** Annual Burn Probability represents the likelihood of a wildfire of any intensity occurring at a given location (pixel) in a single fire season. In a complete assessment of wildfire hazard, wildfire occurrence and spread are simulated in order to characterize how temporal variability in weather and spatial variability in fuel, topography, and ignition density influence wildfire likelihood across a landscape. In such cases, the hazard assessment includes modeling of burn probability, which quantifies the likelihood that a wildfire will burn a given point (a single grid cell or pixel) during a specified period of time. Burn probability for fire management planning applications in this case is reported on an annual basis - the probability of burning during a single fire season.

**Data Resolution:** 30m Raster

**Data Units:** Probability, 0 to 1

**Creation Method:** Annual Burn Probability was produced by Pyrologix LLC, a wildfire threat assessment research firm, as part of a spatial wildfire hazard assessment across all land ownerships for the state of California. The ongoing work generally follows the framework outlined in Scott and Thompson (2013), with custom methods and significant improvements developed by Pyrologix. The project generally consists of three components: fuelscape calibration and updates, wildfire hazard assessment, and risk assessment. It utilizes a combination of wildfire models and custom tools, including the FSim large wildfire simulator (Finney et al., 2011). To date, this work has resulted in a wide variety of spatial data layers related to wildfire hazard and risk, including Annual Burn Probability, representing conditions prior to the 2020, 2021 and 2022 fire seasons. Work to date has been funded by the USDA Forest Service Region 5, the California Energy Commission, and the USDI Bureau of Land Management with data contributions from CAL FIRE.

For this project, the USFS modeling system called FSim is used to quantify annual wildfire likelihood across California. The model is parameterized using spatial datasets of historical weather, fire occurrence, fuels, weather, and topography in order to simulate thousands of fire-years on a landscape. Annual Burn Probability is calculated from these simulations using a Monte Carlo approach to make a spatially resolved estimate of the contemporary annual likelihood of wildfire across the landscape. For more information on FSim or the wildfire hazard modeling being performed by Pyrologix, please see Volger et al., 2021.

**Data Source:** Pyrologix, LLC

**File Name:** AnnualBurnProbability2022.tif

---

## PROBABILITY OF HIGH FIRE SEVERITY

**Tier:** 1

**Data Vintage:** 2022

**Metric Definition and Relevance:** These metrics depicts the probability of high severity fire as constructed by Pyrologix LLC. This operational-control probability raster indicates the probability that the headfire flame length in each pixel will exceed 8 foot flame lengths, the threshold that defines fires that would exceed manual control.

**Data Resolution:** 30m Raster

**Data Units:** Probability, 0 to 1

**Creation Method:** Probability of High Fire Severity (defined as >8 ft) was produced by Pyrologix LLC, a wildfire threat assessment research firm, as part of a spatial wildfire hazard assessment across all land ownerships for the state of California. The ongoing work generally follows the framework outlined in Scott and Thompson (2013), with custom methods and significant improvements developed by Pyrologix. The project generally consists of three components: fuelscape calibration and updates, wildfire hazard assessment, and risk assessment. To date, this work has resulted in a wide variety of spatial data layers related to wildfire hazard and risk, including operational control probabilities based on conditions prior to the 2020, 2021 and 2022 fire seasons. Work to date has been funded by the USDA Forest Service Region 5, the California Energy Commission, and the USDI Bureau of Land Management with data contributions from CAL FIRE. Please reference the Pyrologix 2021 project report (Volger et al., 2021) for more information.

Pyrologix uses the Wildfire Exposure Simulation Tool (WildEST) to develop this data layer, a deterministic wildfire modeling tool that integrates variable weather input variables and weights them based on how they will likely be realized on the landscape. WildEST is more robust than the stochastic intensity values developed with FSim. This is especially true in low wildfire occurrence areas where predicted intensity values from FSim are reliant on a very small sample size of potential weather variables.

**Data Source:** Pyrologix, LLC

**File Name:** ProbabilityHighFireSev\_2022.tif

## FOREST AND SHRUBLAND RESILIENCE

Forest and shrubland resilience is the ability of forest and shrubland vegetation and structure to remain a forest or shrubland in the face of disturbance (e.g., fire, forest management, climate change, etc.). The Forest and Shrubland Resilience Pillar evaluates forest and shrubland vegetation composition and structure to determine its alignment with desired disturbance dynamics and within tolerances of current and future biophysical conditions when considering changes due to climate change. The last 100 years of forest and shrubland management, combined with changing climates, have resulted in forest and shrubland structure and composition which are not resilient to contemporary disturbances. Forest or shrubland structure and composition are one of the few elements of a wildland that management can modify through treatments to improve conditions.

**DESIRED OUTCOME:** Vegetation composition and structure align with topography, desired disturbance dynamics, and landscape conditions, and are adapted to climate change.

## STRUCTURE

Forest or shrubland structure is the spatial distribution of vegetation (live and dead) both vertically and horizontally on the landscape. Prior to European settlement, forests in the Central Coast Region were characterized by heterogeneous spatial patterns replete with individual large trees, gaps, and tree clumps of various sizes – patterns that were shaped by recurrent fire and other disturbances. After a century-plus of fire exclusion, timber harvesting, agricultural development, urbanization, and other land-use practices, the predominant trend across Californian landscapes is that they have become less resilient to natural and human-caused disturbances. In many cases some sort of restoration treatment may be necessary to reverse these trends.

---

### DENSITY – LARGE TREES

**Tier:** 2

**Data Vintage:** 2022

**Metric Definition and Relevance:** Large trees are important to forest managers as they have a greater likelihood of survival from fire, provide sources of seed stock, wildlife habitat, and contribute to other critical processes like carbon storage and nutrient cycling. Large trees are often the focus of management in order to protect existing ones and to foster future ones. In consultation with National Forests, “large trees” have been designated as greater than 30” dbh.

**Data Resolution:** 30m Raster

**Data Units:** Percent live trees per pixel

**Creation Method:** To determine the cutoff for large trees, we used the allometric equations for the Central Coast region (Table 2) that predict diameter at breast height (DBH, breast height - 4.5 ft) as a function of height (HT). Based on these allometric equations, the mean height for large trees, defined as DBH > 30”, is 83.5 ft (Table 3).

| <b>Table 3.</b> Predicted DBH cut-offs as a function of height for trees in the Central Coast Region. Results based on allometric equations reported in Table 2. RMSE = root mean square estimate of the predicted value. |                    |                  |
|---|--------------------|------------------|
| <b>Predicted DBH (in)</b>   | <b>Height (ft)</b> | <b>RMSE (in)</b> |
| 1   | 2.5                | 1.5              |
| 6   | 14.5               | 2.4              |
| 11  | 27                 | 5.3              |
| 24  | 64.5               | 6.8              |
| 30  | 83.5               | 7.3              |

Block statistics were run on California Forest Observatory (CFO) canopy height pixels greater than or equal to 83.5’ (25m) with 3x3 window to calculate the sum for input cells within a 30m rectangular neighborhood. This assigned number of pixels per 30m (900m<sup>2</sup>) cell. Resultant values of 1 through 9 were converted to percent. All background values were calculated to equal 0, meaning 0% large tree existence.

**Data Source:** California Forest Observatory (Salo Sciences), 2020

**File Name:** LargeTreeDensity\_2022.tif

---

## CANOPY LAYER COUNT

**Tier:** 1

**Data Vintage:** Summer 2020

**Metric Definition and Relevance:**

This layer represents the number of distinct vertical canopy layers of trees. Vertical layer count is a proxy for leaf area index, and maps canopy complexity. Since LANDFIRE doesn't support a NoData value, all NoData pixels in canopy fuel metrics were set to 0 in the Landscape files. (e.g., canopy cover was set to 0 in all NoData locations). Topographic data and surface fuel model remain unaltered.

**Data Resolution:** 10m Raster

**Data Units:** Count

**Creation Method:** Each forest structure metric was derived directly from airborne lidar data, hosted by the USGS 3D Elevation Program. However, these data are only available for a small fraction of California's 423,970 km<sup>2</sup> area. To overcome this, we trained deep learning models—a form of pattern recognition—to identify these forest structure patterns in satellite imagery, then mapped each metric statewide.

These algorithms are of the U-net family of neural network architectures that perform pixel-wise regression and classification tasks. The satellite data includes imagery from Sentinel-1 C-band radar sensors and Sentinel-2 multispectral sensors at 10 m spatial resolution, collected in Fall 2019. Future versions will include imagery from PlanetScope multispectral sensors at 3 m resolution.

Downloaded from [California Forest Observatory - Organizations - WIFIRE Commons Data Catalog \(sdsc.edu\)](https://forestobservatory.com/about.html#about). For more information, go to <https://forestobservatory.com/about.html#about>

**Data Source:** California Forest Observatory (Salo Sciences), 2020

**File Name:** CFO\_CanopyLayerCount2020Summer.tif

---

## CANOPY VEG HEIGHT

**Tier:** 1

**Data Vintage:** Summer 2020

**Metric Definition and Relevance:** This layer represents distance between the ground and the top of the canopy. Canopy height is a proxy for aboveground biomass and the amount of foliage that may be consumed in a canopy fire. Since LANDFIRE doesn't support a NoData value, all NoData pixels in canopy fuel metrics were set to 0 in the Landscape files. (e.g., canopy cover was set to 0 in all NoData locations). Topographic data and surface fuel model remain unaltered.

**Data Resolution:** 10m Raster

**Data Units:** meters, min 0 - max 80; each pixel value represents the average height above ground for vegetation within that pixel

**Creation Method:** Each forest structure metric was derived directly from airborne lidar data, hosted by the USGS 3D Elevation Program. However, these data are only available for a small fraction of California's 423,970 km<sup>2</sup> area. To overcome this, we trained deep learning models—a form of pattern recognition—to identify these forest structure patterns in satellite imagery, then mapped each metric statewide.

These algorithms are of the U-net family of neural network architectures that perform pixel-wise regression and classification tasks. The satellite data includes imagery from Sentinel-1 C-band radar sensors and Sentinel-2 multispectral sensors at 10 m spatial resolution, collected in Fall 2019. Future versions will include imagery from PlanetScope multispectral sensors at 3 m resolution.

Downloaded from [California Forest Observatory - Organizations - WIFIRE Commons Data Catalog \(sdsc.edu\)](https://forestobservatory.com/about.html#about). For more information, go to <https://forestobservatory.com/about.html#about>

**Data Source:** California Forest Observatory (Salo Sciences), 2020

**File Name:** CFO\_CanopyHeight2020Summer.tif

---

## CANOPY VEG COVER

**Tier:** 1

**Data Vintage:** Summer 2020

**Metric Definition and Relevance:** This layer represents horizontal cover fraction occupied by tree canopies. Maps community type & fire regime, as well as available habitat for tree-dwelling species.

**Data Resolution:** 10m Raster

**Data Units:** Canopy cover is a 0-100% cover fraction and may be more precisely described as "canopy density." It calculates the proportion of all lidar returns  $\geq 5$ m divided by the total number of returns in that grid cell. This, therefore, does not include all vegetation, but instead describes the density of vegetation in the canopy vertical stratum (veg 5m and taller).

**Creation Method:** Each forest structure metric was derived directly from airborne lidar data, hosted by the USGS 3D Elevation Program. However, these data are only available for a small fraction of California's 423,970 km<sup>2</sup> area. To overcome this, we trained deep learning models—a form of pattern recognition—to identify these forest structure patterns in satellite imagery, then mapped each metric statewide.

These algorithms are of the U-net family of neural network architectures that perform pixel-wise regression and classification tasks. The satellite data includes imagery from Sentinel-1 C-band radar sensors and Sentinel-2 multispectral sensors at 10 m spatial resolution, collected in Fall 2019. Future versions will include imagery from PlanetScope multispectral sensors at 3 m resolution.

Downloaded from [California Forest Observatory - Organizations - WIFIRE Commons Data Catalog \(sdsc.edu\)](https://forestobservatory.com/about.html#about). For more information, go to <https://forestobservatory.com/about.html#about>

**Data Source:** California Forest Observatory (Salo Sciences), 2020

**File Name:** CFO\_CanopyCover2020Summer.tif

## COMPOSITION

The composition of a forest is a reference to the biodiversity of the landscape; this includes a diversity of vegetation species, types (e.g., trees, shrubs, forbs, etc.), and distribution. Tree species composition affects many aspects of forest dynamics and function. A diversity of tree and shrub species can confer greater resilience to climate change and beetle outbreaks. The vegetation composition also affects fire dynamics, water reliability, carbon pools and sequestration, and economic diversity pillars. Since European settlement and the adoption of fire suppression and logging, forests of the Central Coast Region shifted to increased dominance of shade-tolerant and fire-intolerant species like white fir and red fir, incense cedar, Douglas fir, and tanoak. Other species like ponderosa pine, Jeffrey pine, sugar pine, and black oak, which are more shade-intolerant and fire-tolerant, declined in coverage. With increasingly larger and higher-severity fires occurring, forest-cover loss may be significant and shrub cover will increase.

---

## TREE COVER

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Total tree cover as measured by the fractional non-overlapping absolute tree cover, viewed vertically. Provides a first order measure of vegetation type when combined with parallel observations of shrub and herbaceous cover. Data from the National Land Cover Database (NLCD) are used for training, and NLCD definitions for cover (for example, the distinction between tree vs shrub) are expected to be similar in the CECS data sets.

**Data Resolution:** 30m Raster

**Data Units:** Fractional non-overlapping absolute cover; continuous variable from 0 to 1.

**Creation Method:** Machine learning (Random Forest) using the National Land Cover Database for training and Landsat observations as predictors. See <https://doi.org/10.1029/2021AV000654> for further information.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** VegCover\_Tree\_2021.tif

---

## SHRUB COVER

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Total shrub cover as measured by the fractional non-overlapping absolute shrub cover, viewed vertically. Provides a first order measure of vegetation type when combined with parallel observations of tree and herbaceous cover. Data from the National Land Cover Database (NLCD) are used for training, and NLCD definitions for cover (for example, the distinction between tree vs shrub) are expected to be similar in the CECS data sets.

**Data Resolution:** 30m Raster

**Data Units:** Fractional non-overlapping absolute cover; continuous variable from 0 to 1.

**Creation Method:** Machine learning (Random Forest) using the National Land Cover Database for training and Landsat observations as predictors. See <https://doi.org/10.1029/2021AV000654> for further information.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** VegCover\_Shrub\_2021.tif

---

## HERBACEOUS COVER

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Total herbaceous cover as measured by the fractional non-overlapping absolute herbaceous cover, viewed vertically. Provides a first order measure of vegetation type when combined with parallel observations of tree and herbaceous cover. Data from the National Land Cover Database (NLCD) are used for training, and NLCD definitions for cover (for example, the distinction between tree vs shrub) are expected to be similar in the CECS data sets.

**Data Resolution:** 30m Raster

**Data Units:** Fractional non-overlapping absolute cover; continuous variable from 0 to 1.

**Creation Method:** Machine learning (Random Forest) using the National Land Cover Database for training and Landsat observations as predictors. See <https://doi.org/10.1029/2021AV000654> for further information.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** VegCover\_Herb\_2021.tif

---

## SERAL STAGE

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** The seral stages are categories that represent the developmental progression of forest ecosystems from initial establishment or following a stand replacing event (e.g., high severity fire) to a forest dominated by trees in the upper age classes for a given forest type. Late seral forests are also often characterized by multiple ages of forest trees and dead and dying trees in some form of equilibrium. Seral conditions across landscapes were highly variable prior to major European settlement in the western US. These patterns were highly attuned to dominant disturbance regimes and the multi-scaled variability in environmental conditions across topographic and climatic gradients. These patterns helped to reinforce fire regimes dominated by low- to moderate-severity fire across much of the region and provided for multiple habitat requirements for a wide variety of species.

This metric contains three related data layers. The first is an assignment to each 30 meter pixel of the seral stage it is currently in, either early, mid, or late seral stage. The other two layers represent the proportion of a HUC 12 watershed that is in 1) early seral stage or 2) late seral stage.

**Data Resolution:** 30m Raster, HUC 12 watersheds

**Data Units:** Integer 1 - 3, continuous variable 0-1

**Creation Method:** The FVEG data, used in characterizing vegetation and habitat conditions for a number of metrics in this kit, contain data on tree size ([see FVEG discussion above](#)). Seral stages for forested lands are binned into one of three categories of tree size (Early, Mid, Late) and those are defined by tree diameter, per the CWHR system.



| Size Class        | Size (inches DBH) | Seral Stage |
|-------------------|-------------------|-------------|
| 1 Seedling        | less than 1       | Early (1)   |
| 2 Sapling         | 1 – 6             | Early (1)   |
| 3 Pole            | 6 – 11            | Mid (2)     |
| 4 Small           | 11 – 24           | Mid (2)     |
| 5 Medium to Large | 24+               | Late (3)    |
| 6 Multi-storied   | 36 – 48           | Late (3)    |

Late Seral conditions have been lumped into a single classification (24” and up).

The first layer provided here assigns a early, mid, or late seral value to each cell based on dominant tree size in the canopy. The second and third data layer provided identify the proportion of the HUC12-scale (typically 10,000-30,000 acres in size) that is either early seral forest or late seral forest, respectively. These patterns can be highly variable at finer-scales so we used a HUC 12 watershed as the unit for expressing relative abundance. For each HUC12, the proportion of the watershed covered by the evaluated seral stage has been calculated.

**Data Source:** FVEG

**File Name:** SeralStage\_EML.tif; early\_SeralStage\_prop.tif; late\_SeralStage\_prop.tif

## DISTURBANCE

Central Coast forests evolved with a suite of frequent disturbances: wildfires (both from lightning and burning by indigenous people), bark beetle-caused mortality, drought-caused mortality, avalanches, landslides, and windthrow, all of which created forest heterogeneity across the landscape. This heterogeneity included variations in surface and ladder fuels, which moderated fire behavior and spread. The variations in stand density and forest opening also served as critical habitats for wildlife. Forested areas are now more homogeneous due to lack of disturbance. The lack of disturbance is evident in the forest structure.

---

## CUMULATIVE TREE COVER LOSS

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** The cumulative loss of tree cover over a 30-year period (1992-2021). Tree cover loss reflects fires, harvest/management and dieoff. Only disturbances that are sufficient to trigger the Continuous Change Detection and Classification algorithm are included; low-level, diffuse dieoff is likely missed.

**Data Resolution:** 30m Raster

**Data Units:** Cumulative fractional non-overlapping absolute tree cover loss, where tree cover is a continuous variable from 0 to 1. Cumulative loss can exceed 1 in cases with multiple disturbances.

**Creation Method:** Vegetation disturbances were identified over the Landsat TM/ETM+/OLI era using the Continuous Change Detection and Classification algorithm (CCDC). The corresponding annual change in tree cover was determined with machine learning (Random Forest) using the National Land Cover Database for training and Landsat/CCDC observations as predictors; this produced a ~35-year stack of rasters that identified the locations and severity of tree cover loss. This stack was then summed for 1992-2021 to calculate the cumulative tree cover loss over a 30-year period. See <https://doi.org/10.1029/2021AV000654> for further information.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** DistHist\_Severe\_Tree\_19922021.tif

---

## CUMULATIVE SHRUB COVER LOST

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** The cumulative loss of shrub cover over a 30-year period (1992-2021). Shrub cover loss reflects fires, harvest/management and dieoff. Only disturbances that are sufficient to trigger the Continuous Change Detection and Classification algorithm are included; low-level, diffuse dieoff is likely missed.

**Data Resolution:** 30m Raster

**Data Units:** Cumulative fractional non-overlapping absolute shrub cover loss, where shrub cover is a continuous variable from 0 to 1. Cumulative loss can exceed 1 in cases with multiple disturbances.

**Creation Method:** Vegetation disturbances were identified over the Landsat TM/ETM+/OLI era using the Continuous Change Detection and Classification algorithm (CCDC). The corresponding annual change in shrub cover was determined with machine learning (Random Forest) using the National Land Cover Database for training and Landsat/CCDC observations as predictors; this produced a ~35-year stack of rasters that identified the locations and severity of shrub cover loss. This stack was then summed for 1992-2021 to calculate the cumulative tree cover loss over a 30-year period. See <https://doi.org/10.1029/2021AV000654> for further information.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** DistHist\_Severe\_Shrub\_19922021.tif

---

## RISK OF TREE DIEOFF DURING DROUGHT

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** A quantitative continuous variable that reflects the risk of tree dieoff during a significant drought period (SPI48 drought = -2).

**Data Resolution:** 30m Raster

**Data Units:** This is a dimensionless index that ranges from 0 to ~20000. Low values indicate minimal or no risk of tree dieoff during drought, either or both because there are few trees in the pixel and/or there is ample local moisture even during periods of extreme precipitation shortfall. High values indicate significant risk of tree dieoff during drought, as a result of both a high density of trees at the site and likelihood of extreme local moisture shortfall.

**Creation Method:** Calculated by combining information on the local moisture balance and tree density. Local moisture balance was calculated as the ratio of Annual Evapotranspiration with the canopy observed in 2021 to Precipitation during a SPI 48 drought = -2 based on local P observations during 1991-2020. This ratio quantifies the local moisture deficit/surplus that would be expected during a 48-month period with precipitation that is 2 standard deviations below the local 30 year Normal. Tree cover was determined from Landsat. See <https://doi.org/10.1038/s41561-019-0388-5> for further information.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

---

POTENTIAL CLIMATE REFUGIA -BASELINE (HISTORICAL) CONDITIONS

Tier: 3

Data Vintage: 2016

**Metric Definition and Relevance:** This raster dataset represents habitat types (natural vegetation communities) and their distribution across the array of climate conditions that each separate habitat type is found in under the baseline climate conditions. A 2015 map of the state’s natural vegetation compiled from multiple sources was classified to the National Vegetation Classification Standard’s mid-level classification, called “Macrogroup”. Thirty one natural vegetation macrogroups are identified in the map, covering 99.87% of the state’s natural terrestrial vegetation, and occupying 353,271 km<sup>2</sup>.

This serves as the foundation from which habitat types will be exposed to predicted changes in climate. Data are arrayed across 0 to 1 in terms of their exposure to current climate conditions. This data layer provides a baseline of vegetation adapted to “historic” conditions; i.e. climate conditions from the recent past; 1980-2010.

**Data Resolution:** 270m Raster

**Data Units:** 0- 1. Low values indicate higher resilience to threats. High values indicate significant exposure to climate change. -1 represents ‘non analog’ areas, i.e. locations that are outside the historic climate envelope of a given vegetation type.

**Creation Method:** The vegetation climate exposure analysis takes advantage of the 2015 vegetation map compiled for California by CALFIRE. Each Macrogroup (MG) was analyzed to determine which California habitats and associated dominant plant species make up its definition. California habitats are defined by the California Department of Fish and Wildlife (CDFW) through their California Wildlife Habitat Relationship (WHR) models<sup>9</sup>. WHR types are made up of plant species, such as the dominant trees, shrubs, and smaller plants. CDFW experts determined which WHR types correspond to each individual macrogroup; this cross-walk was used to develop a list of the dominant plant species that comprise each macrogroup.

The climate space occupied by each distinct vegetation macrogroup (largely equivalent to a CWHR habitat type) from the current time period was identified. This was done by using the points for each type and applying a kernel density estimator on a 2-d surface composed of the first two principal components of the climate conditions. The result is a smoothed continuous point density surface, showing the prevalence of each vegetation type across the range of sampled climatic conditions. This surface was partitioned by fitting contour lines so that they enclose a proportion of the original points from the current time period. Contours were calculated at 5% increments. For example the innermost 5% contour line encloses the 5% of pixels for the given vegetation type which are at the core of the climate space for that type, as determined by its density in the climate space. Cells further away from the dense central core, are considered to be more marginal in the vegetation type’s distribution. The outer contours are fit to enclose the 95-99% of climatically marginal points, with the last 1% of cells (beyond the 99% contour) being the most marginal. In addition, if a cell lies outside the space defined by the 99% contour of any vegetation type, it is considered to be “non-analog,” which means that it experiences climatic conditions outside of the conditions where we have a good sample in the initial time period. Excluded from this assessment are non-vegetated types such as snow, open water, and ice; and non-natural landcover types mapped as vineyards, tilled earth, orchards and Urban.

For more information on methods for the development of these climate refugia data see:

Thorne et al. 2015

Thorne et al. 2016

Thorne et al. 2017

Thorne et al. 2020

**Data Source:** Information Center for the Environment, UC Davis

**File Name:** hst8110.tif

---

## POTENTIAL CLIMATE REFUGIA - UNDER MODELED CLIMATE CHANGE (MIROC MODEL - HOTTER AND DRIER)

**Tier:** 3

**Data Vintage:** 2016

**Metric Definition and Relevance:** This raster dataset represents habitat types (CWHR habitat classes) and their predicted exposure to climate stress across the array of predicted climate conditions (separate layers for early (2010 - 2039), mid (2040-2069), and late century (2070-2099)) for all habitat types in comparison to the baseline climate conditions. This serves as the foundation from which habitat types will be exposed to predicted changes in climate. Data are arrayed across 0 to 1 in terms of their exposure to current climate conditions. These three data layers can be used to help land managers allocate limited resources for climate-adaptive field work by providing a view of climate risk that varies across the lands they manage.

The Climate Change Model used in this analysis is the Miroc Earth System Model. This ESM, named “MIROC-ESM”, is based on a global climate model MIROC (Model for Interdisciplinary Research on Climate) which has been cooperatively developed by researchers in Japan and others. This model suggests a hotter and drier future. The emission scenario used is the RCP 8.5, which represents a range of warming statewide from 1.99 to 4.56°C and between a 24.8% decrease in precipitation and a 22.9% increase, respectively.

**Data Resolution:** 270m Raster

**Data Units:** 0- 1. Low values indicate higher resilience to threats. High values indicate significant exposure to climate change. -1 represents ‘non analog’ areas, i.e. locations that are outside the historic climate envelope of a given vegetation type.

**Creation Method:** The vegetation climate exposure analysis takes advantage of the 2015 vegetation map compiled for California, which is described above. The vegetation climate exposure model is implemented in the R programming language, and takes the vegetation and climate raster files as the primary input data. The values of the climate raster files were randomly sampled at 100,000 points on the landscape, which were used to fit a statistical model characterizing the relationship between the variables both in the current time and for the modeled future data.

At each of these 100,000 points, 9 hydro-climatic variables were sampled to characterize the range and variation of conditions in the study region. These variables were: annual mean minimum temperature (Tmin), annual mean maximum temperature (Tmax), annual precipitation (PPT), actual evapotranspiration (AET), potential evapotranspiration (PET), climatic water deficit (CWD), snowpack depth on April 1st, runoff, and recharge. The variation between these variables was modeled using a principal component analysis<sup>21</sup> (PCA) to identify the

dominant components of variation. The top-two principal components axes, representing about 79% of the variability across the four climate projections, were extracted as a two-dimensional space, and are portrayed as the axes for the PCA plots shown in each macrogroup chapter below. This was done to simplify the representation of the climate space, while maintaining the most important information on the variables to be associated with the observed vegetation distributions.

The climate space occupied by each distinct macrogroup from the current time period was identified. This was done by using the points for each type and applying a kernel density estimator on a 2-d surface composed of the first two principal components of the climate conditions. The result is a smoothed continuous point density surface, showing the prevalence of each vegetation type across the range of sampled climatic conditions. This surface was partitioned by fitting contour lines so that they enclose a proportion of the original points from the current time period. Contours were calculated at 5% increments. For example the innermost 5% contour line encloses the 5% of pixels for the given vegetation type which are at the core of the climate space for that type, as determined by its density in the climate space. Cells further away from the dense central core, are considered to be more marginal in the vegetation type's distribution. The outer contours are fit to enclose the 95-99% of climatically marginal points, with the last 1% of cells (beyond the 99% contour) being the most marginal. In addition, if a cell lies outside the space defined by the 99% contour of any vegetation type, it is considered to be "non-analog," which means that it experiences climatic conditions outside of the conditions where we have a good sample in the initial time period. As a result, the status of that point is uncertain. There are occasionally a few extreme points which appear to be far outside the general distribution for the type. These may be due to misclassified vegetation types in the source data, microclimatic conditions not captured by the climate data, historic anomalies in long-lived species, etc.

Climate exposure is the level of climate change expected in the areas where each macrogroup is dominating. This report uses the term "vegetation climate exposure analysis" to describe the following analysis which was conducted on each macrogroup. The vegetation climate exposure analysis is calculated using the mapped extent of each macrogroup. Every grid cell of each macrogroup was ranked as to its level of exposure, relative to the entire area of that macrogroup. This was done for the current time, and used to define the common climate found for each macrogroup. Once each type's "climate envelope" was defined, we then assessed how much every grid cell changed under various future climate projections. This allowed a measure of the vegetation stress, or climate exposure. The area extent of each macrogroup that will be lost from the most commonly occurring climate conditions ( $\leq 80\%$ ) and the area that will fall into current marginal, or stressed, climate conditions ( $> 95\%$ ) or outside the current climate conditions was calculated. This approach is particularly useful for resource managers, who often are constrained to work in specified areas, and need estimates of what areas within their jurisdiction are likely to be highly stressed, and what areas are likely to be less stressed, in effect climate refuge areas.

To consider how refugial conditions from a range of stressors can inform conservation planning and management, the authors integrated metrics of refugial capacity across different domains, which are defined as social, ecological, or physical drivers, processes, or cycles that influence landscape structure, function, or composition. To persist in the California landscape, species and ecosystems may need refugia from shifting climatic conditions, including extremely hot summers and prolonged droughts, but non-climate stressors can also affect conservation outcomes. In this landscape, changes in fire frequency can be a significant stressor affecting plant community structure and persistence. Anthropogenic features that modify hydrologic flows alter the ability of watersheds to sustain functional habitats. And finally, protected areas are often designed to mitigate the impacts of anthropogenic activities; however, recreational activities may alter the refugial capacity of the protected land, affecting the ability of the landscape to sustain species and their habitats. We combined these individual metrics to assess landscape level refugial capacity.

Sites with high refugial capacity (super-refugia sites) have, on average, 30% fewer extremely warm summers, 20% fewer fire events, 10% less exposure to altered river channels and riparian areas, and 50% fewer recreational trails than the surrounding landscape. Our results suggest that super-refugia sites (~8,200 km<sup>2</sup>) for some natural communities are underrepresented in the existing protected area network, a finding that can inform efforts to expand protected areas.

For more information on methods for the development of these climate refugia data see:

Thorne et al. 2015

Thorne et al. 2016

Thorne et al. 2017

Thorne et al. 2020

**Data Source:** Information Center for the Environment, UC Davis

**File Name:** miroc\_85\_1039.tif; miroc\_85\_4069.tif; miroc\_85\_7099.tif

---

## POTENTIAL CLIMATE REFUGIA - COMBINED MODELED CLIMATE CHANGE (MIROC MODEL - (HOTTER AND DRIER) AND CNRM-CM5 (WETTER AND WARMER))

**Tier:** 3

**Data Vintage:** 2016

**Metric Definition and Relevance:** This raster dataset represents habitat types (Macro Veg Type, largely equivalent to CWHR habitat classes) and their predicted exposure to climate stress across the array of predicted climate conditions (separate layers for early (2010 - 2039), mid (2040-2069), and late century (2070-2099)) for all habitat types in comparison to the baseline climate conditions. This serves as the foundation from which habitat types will be exposed to predicted changes in climate. Data are arrayed across 0 to 1 in terms of their exposure to current climate conditions. These three data layers can be used to help land managers allocate limited resources for climate-adaptive field work by providing a view of climate risk that varies across the lands they manage.

This analysis uses both the Miroc Earth System Model and the CNRM-CM5. CNRM-CM5 is an Earth system model designed to run climate simulations. It consists of several existing models designed independently and coupled through the OASIS software. Both were used under the RCP 8.5 emission scenario given that this is more likely under current emission levels.

This data layer is provided as a summary of likely exposure results. **Exposure Scores:**

- 1 = Refugia: CNRM-CM5 only (CNRM exposure values < 80%)
- 2 = Refugia: MIROC-ESM only (MIROC exposure values < 80%)
- 3 = Refugia Consensus (both models agree exposure values < 80%)
- **8 = High Exposure (both models agree exposure values >95%)**
- **9 = Very High Exposure (both models agree exposure values >99%)**

**Data Resolution:** 270m Raster

**Data Units:** 0, 1, 2, 3, 8, 9 Low values indicate higher resilience to threats. High values indicate significant exposure to climate change. -1 represents 'non analog' areas, i.e. locations that are outside the historic climate envelope of a given vegetation type.

**Creation Method:** Each dominant species is scored for its sensitivity to, and ability to adapt (adaptive capacity) to climate change. Sensitivity refers to the degree to which changes in climate are thought to directly impact different species. Adaptive capacity refers to estimates of the degree to which different species can use their life history characteristics to moderate impacts from changing climate. These two sets of scores represent the biological attributes of the dominant species in each macrogroup. We scored each of the dominant species comprising each macrogroup, according to life history characteristics defined in attribute tables of the California Manual of Vegetation, and supplemented by information found in the USDA plants database and the Jepson Interchange, a web portal for California plant taxonomy. The scores were combined to generate a single sensitivity and adaptive capacity (S&A) score.

Climate exposure is the level of climate change expected in the areas where each macrogroup is dominating. This report uses the term “vegetation climate exposure analysis” to describe the following analysis which was conducted on each macrogroup. The vegetation climate exposure analysis is calculated using the mapped extent of each macrogroup. Every grid cell of each macrogroup was ranked as to its level of exposure, relative to the entire area of that macrogroup. This was done for the current time, and used to define the common climate found for each macrogroup. Once each type’s “climate envelope” was defined, we then assessed how much every grid cell changed under various future climate projections. This allowed a measure of the vegetation stress, or climate exposure. The area extent of each macrogroup that will be lost from the most commonly occurring climate conditions ( $\leq 80\%$ ) and the area that will fall into current marginal, or stressed, climate conditions ( $> 95\%$ ) or outside the current climate conditions was calculated. This approach is particularly useful for resource managers, who often are constrained to work in specified areas, and need estimates of what areas within their jurisdiction are likely to be highly stressed, and what areas are likely to be less stressed, in effect climate refuge areas.

To consider how refugial conditions from a range of stressors can inform conservation planning and management, the authors integrated metrics of refugial capacity across different domains, which are defined as social, ecological, or physical drivers, processes, or cycles that influence landscape structure, function, or composition. To persist in the California landscape, species and ecosystems may need refugia from shifting climatic conditions, including extremely hot summers and prolonged droughts, but non-climate stressors can also affect conservation outcomes. In this landscape, changes in fire frequency can be a significant stressor affecting plant community structure and persistence. Anthropogenic features that modify hydrologic flows alter the ability of watersheds to sustain functional habitats. And finally, protected areas are often designed to mitigate the impacts of anthropogenic activities; however, recreational activities may alter the refugial capacity of the protected land, affecting the ability of the landscape to sustain species and their habitats. We combined these individual metrics to assess landscape level refugial capacity.

Sites with high refugial capacity (super-refugia sites) have, on average, 30% fewer extremely warm summers, 20% fewer fire events, 10% less exposure to altered river channels and riparian areas, and 50% fewer recreational trails than the surrounding landscape. Our results suggest that super-refugia sites ( $\sim 8,200 \text{ km}^2$ ) for some natural communities are underrepresented in the existing protected area network, a finding that can inform efforts to expand protected areas.

For more information on methods for the development of these climate refugia data see:

Thorne et al. 2015

Thorne et al. 2016

Thorne et al. 2017

Thorne et al. 2020

**Data Source:** Information Center for the Environment, UC Davis

**File Name:** combine85\_all7.tif

## BIODIVERSITY CONSERVATION

The California landscape provides habitat for over 300 species of native vertebrates and thousands of invertebrate species and plants. Management activities over the last century have impacted most species to varying degrees and some have declined significantly in recent decades. Protecting and enhancing native biodiversity has become a management imperative under both federal and state laws and policy. Native plants and animals provide a wide array of benefits to forests and other habitats in California; they help forests recover after a fire, control flooding and soil erosion, cycle nutrients, and are valued by people recreating in forests. Greater species diversity promotes adaptability and helps ecosystems withstand and recover from disturbance, including those caused by climate change. The Biodiversity Conservation pillar focuses on species diversity, critical habitat for focal species and non-native species distribution.

Habitat data to model the likelihood of species presence or absence was derived from the [FVEG WHR data layer](#).

**DESIRED OUTCOME:** The network of native species and ecological communities is sufficiently abundant and distributed across the landscape to support and sustain their full suite of ecological and cultural roles.

## SPECIES DIVERSITY

Species diversity is a function of both the number of different species in the community and their relative abundances. Larger numbers of species and more even abundances of species lead to higher species diversity. Species diversity can be calculated in a variety of ways to represent the type and magnitude of differences among species, their number, and their abundance.

---

## WILDLIFE SPECIES RICHNESS

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** Native species richness is estimated based on high suitability reproductive habitat for a given species. Reproductive habitat is used to represent suitability because it is critical for species persistence and for most native species it has the most limited requirements. If a habitat is identified as high for a given species, it is considered suitable (1), and habitat identified as moderate, low or not suitable, it is considered unsuitable (0). Species richness values are used as a relative measure of biodiversity value; as such, areas with lower species richness based on these criteria may still have high biodiversity value, but not as high as areas with higher richness values. The number of native species per spatial unit (30m pixel) presented as simply the total number; this can be useful for assessing change in number/composition over space. These values are specific to the Central Coast species and footprint for this kit.



**Data Resolution:** 30m Raster

**Data Units:** Number of species

**Creation Method:** Generated using the California Wildlife Habitat Relationships model developed and managed by the California Department of Fish and Wildlife. CWHR habitat values are based on the FVEG vegetation data that has been updated. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction of that species in the California Wildlife Habitat Relationship database.

**Data Source:**

CDFW

CALFIRE

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** wildlife\_species\_richness.tif

---

## THREATENED/ENDANGERED VERTEBRATE SPECIES RICHNESS

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** Native species richness is estimated based on high suitability reproductive habitat for a given species. Reproductive habitat is used to represent suitability because it is critical for species persistence and for most native species it has the most limited requirements. If a habitat is identified as high for a given species, it is considered suitable (1), and habitat identified as moderate, low or not suitable, it is considered unsuitable (0). Species richness values are used as a relative measure of biodiversity value; as such, areas with lower species richness based on these criteria may still have high biodiversity value, but not as high as areas with higher richness values. The total number of federally threatened/endangered native species per spatial unit (30m pixel) can be useful for assessing change in number/composition over space. These values are specific to the Central Coast species and footprint for this kit.

**Data Resolution:** 30m Raster

**Data Units:** Number of species

**Creation Method:** Generated using the California Wildlife Habitat Relationships model developed and managed by the California Department of Fish and Wildlife. CWHR habitat values are based on the FVEG vegetation data that has been updated. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction of that species in the California Wildlife Habitat Relationship database.

Only species classified in the CWHR database as federally endangered, federally threatened, California endangered, or California threatened have been included in the species richness count for this layer.

**Data Source:**

CDFW

CALFIRE

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** t\_e\_species\_richness.tif

---

## FOREST RAPTORS SPECIES RICHNESS

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** Native species richness is estimated based on high suitability reproductive habitat for a given species. Reproductive habitat is used to represent suitability because it is critical for species persistence and for most native species it has the most limited requirements. If a habitat is identified as high for a given species, it is considered suitable (1), and habitat identified as moderate, low or not suitable, it is considered unsuitable (0). Species richness values are used as a relative measure of biodiversity value; as such, areas with lower species richness based on these criteria may still have high biodiversity value, but not as high as areas with higher richness values. The total number of federally threatened/endangered native species per spatial unit (30m pixel) can be useful for assessing change in number/composition over space.

**Data Resolution:** 30m Raster

**Data Units:** Number of species

**Creation Method:** Generated using the California Wildlife Habitat Relationships model developed and managed by the California Department of Fish and Wildlife. CWHR habitat values are based on the FVEG vegetation data that has been updated. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction of that species in the California Wildlife Habitat Relationship database.

Only raptor species that are associated with forest habitats have been included in the species richness count for this layer. The raptors included in this layer are Bald Eagle, California Spotted Owl, Cooper'S Hawk, Great-Horned Owl, Merlin, Northern Goshawk, Northern Spotted Owl, Osprey, Peregrine Falcon, Red-Shouldered Hawk, Red-Tailed Hawk, Screech Owl and Sharp-Shinned Hawk.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** forest\_species\_richness.tif

---

## OPEN HABITAT RAPTORS SPECIES RICHNESS

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** Native species richness is estimated based on high suitability reproductive habitat for a given species. Reproductive habitat is used to represent suitability because it is critical for species

persistence and for most native species it has the most limited requirements. If a habitat is identified as high for a given species, it is considered suitable (1), and habitat identified as moderate, low or not suitable, it is considered unsuitable (0). Species richness values are used as a relative measure of biodiversity value; as such, areas with lower species richness based on these criteria may still have high biodiversity value, but not as high as areas with higher richness values. The total number of federally threatened/endangered native species per spatial unit (30m pixel) can be useful for assessing change in number/composition over space.

**Data Resolution:** 30m Raster

**Data Units:** Number of species

**Creation Method:** Generated using the California Wildlife Habitat Relationships model developed and managed by the California Department of Fish and Wildlife. CWHR habitat values are based on the FVEG vegetation data that has been updated. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction of that species in the California Wildlife Habitat Relationship database.

Only raptor species that are associated with open habitats have been included in the species richness count for this layer. The raptors included in this layer are American Kestrel, Barn Owl, Burrowing Owl, Ferruginous Hawk, Golden Eagle, Long-Eared Owl, Northern Harrier, Prairie Falcon, Rough-Legged Hawk, Short-Eared Owl, Swainson'S Hawk, Turkey Vulture, White-Tailed Kite, California Condor and Great Grey Owl.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** open\_species\_richness.tif

---

## HUMMINGBIRDS SPECIES RICHNESS

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** Native species richness is estimated based on high suitability reproductive habitat for a given species. Reproductive habitat is used to represent suitability because it is critical for species persistence and for most native species it has the most limited requirements. If a habitat is identified as high for a given species, it is considered suitable (1), and habitat identified as moderate, low or not suitable, it is considered unsuitable (0). Species richness values are used as a relative measure of biodiversity value; as such, areas with lower species richness based on these criteria may still have high biodiversity value, but not as high as areas with higher richness values. The total number of federally threatened/endangered native species per spatial unit (30m pixel) can be useful for assessing change in number/composition over space.

**Data Resolution:** 30m Raster

**Data Units:** Number of species

**Creation Method:** Generated using the California Wildlife Habitat Relationships model developed and managed by the California Department of Fish and Wildlife. CWHR habitat values are based on the FVEG vegetation data that has been updated. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction of that species in the California Wildlife Habitat Relationship database.

Only hummingbird species have been included in the species richness count for this layer.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** hummingbirds\_species\_richness.tif

## FOCAL SPECIES

For specified species listed below within the Focal Species element section of the Biodiversity Conservation pillar, the species should be considered as *Species of Interest*. It is important for the readers to understand, the listed species are not exhaustive, may be an Endangered Species Act (ESA) species, or considered Sensitive Species as they pertain to forest planning. These species are identified based on their sensitivity to impacts from restoration thinning, prescribed fire, and wildfire. The two wildlife species are California spotted owl and fisher. Black oak is an important species for wildlife as well as for tribes.

---

### CALIFORNIA SPOTTED OWL

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** California spotted owl is distributed in the central coast region from Monterey County to Santa Barbara County and inhabits elevations ranging from 1,000 to over 7,000 feet. It is a Region 5 Forest Service “Sensitive Species” and a “Management Indicator Species” (representing late seral closed canopy coniferous forest). In 2023, the USFWS issued a 12-month finding on a petition to list the California spotted owl under the Endangered Species Act and determined listing to be not warranted at this time (USDI Fish and Wildlife Service 2023). The species is declining throughout much of its range and faces continued threats due to wildfire, habitat loss, and competition from barred owls. A conservation assessment for California spotted owl was conducted in 2017 (Gutiérrez, Manley, and Stine 2017). This was followed by the development of a conservation strategy to guide habitat management on National Forest System Lands (USDA Forest Service 2019). The conservation strategy for the California spotted owl throughout its range, including the Central Coast, aims to balance the need to conserve essential habitat elements around sites occupied by California spotted owls, while simultaneously restoring resilient forest conditions at the landscape scale (USDA Forest Service 2019).

The USDA Forest Service designates a 300-acre protected activity center (PAC) around each known nesting area or activity center. PACs are a USFS land allocation designed to protect and maintain high-quality California spotted owl nesting and roosting habitat around active sites.

The map associated with this data layer includes the southern extent of the Northern Spotted Owl (Marin County).

**Data Resolution:** 30m Raster

**Data Units:** Binary, 0 (Low Suitability), 1 (High Suitability)

**Creation Method:** CWHR classifications are based on a combination of FVEG canopy cover, FVEG size class and vegetation data. The vegetation data includes a variety of tree, shrub, grassland, and water dominated habitats. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed suitable for the reproduction of that species in the California Wildlife Habitat Relationship database. Habitat that meets the following criteria is considered suitable:

- CWHR size and density of 4D, 4M within CWHR vegetation types of DFR, MHC, MHW, MRI, PPN, RFR, SMC, WFR
- CWHR size and density of 5D, 5M, 6 within CWHR vegetation types of DFR, EPN, JPN, LPN, MHC, MHW, MRI, PPN, RFR, SMC, WFR

CWHR high suitability values have been used to create separate data layers which identify suitable nesting and suitable foraging habitat. These data have been combined to create the identified “suitable habitat” layers.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

Conservation Strategy for the California Spotted Owl in the Sierra Nevada, US Forest Service, 2019

**File Name:** California\_Spotted\_Owl\_suitable\_habitat.tif

---

## MOUNTAIN LION

**Tier:** 2

**Data vintage:** 2014

**Metric Definition and Relevance:** This layer shows highly suitable habitats for the reproduction and feeding of Mountain lion (*Puma concolor*).

**Data Resolution:** 30m Raster

**Data Units:** Binary, 0 (not suitable) and 1 (suitable)

**Creation Method:** CWHR classifications are based on a combination of FVEG canopy cover, FVEG size class and vegetation data. The vegetation data includes a variety of tree, shrub, grassland, and water dominated habitats. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction or feeding of that species in the California Wildlife Habitat Relationship database.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** Mountain\_Lion\_suitable\_habitat.tif

---

## CALIFORNIA RED-LEGGED FROG

**Tier:** 1

**Data Vintage:** 2001

**Metric Definition and Relevance:** This dataset represents a species habitat distribution map for California Red-legged Frog (*Rana draytonii*) within the conterminous United States (CONUS) based on 2001 ground conditions.

**Data Resolution:** 30m Raster

**Data Units:** Binary layer, 1 represents current habitat

**Creation Method:** This Gap Analysis Project (GAP) habitat map is a prediction of the spatial distribution of suitable environmental and land cover conditions within the United States for the species. Mapped areas represent places where the environment is suitable for the species to occur (i.e. suitable to support one or more life history requirements for breeding, resting, or foraging), while areas not included in the map are those predicted to be unsuitable for the species. While the actual distributions of many species are likely to be habitat limited, suitable habitat will not always be occupied because of population dynamics and species interactions. Furthermore, these maps correspond to midscale characterizations of landscapes, but individual animals may deem areas to be unsuitable because of presence or absence of fine-scale features and characteristics that are not represented in our models (e.g. snags, vernal pools, shrubby undergrowth). These maps are intended to be used at a 1:100,000 or smaller map scale.

This habitat map is created using a deductive model to predict areas suitable for occupation within a species range. The deductive habitat models are built by compiling information on the species' habitat associations and entering it into a relational database. Information is compiled from the best available characterizations of the species' habitat, which included species accounts in books and databases, primary peer-reviewed literature. The literature references for each species are included in the "Species Habitat Model Report" and "Machine Readable Habitat Database Parameters" files attached to each habitat map item in the ScienceBase repository. The compiled habitat information is used by a biologist to determine which of the ecological systems and land use classes represented in the National Gap Analysis Project's (GAP) Land Cover Map Ver. 1.0 the species is associated with.

The maps are generated using a python script that queries the model parameters in the database; reclassifies the GAP Land Cover Ver 1.0 and ancillary data layers within the species' range; and combines the reclassified layers to produce the final 30m resolution habitat map. Map output is, therefore, not only a reflection of the ecological systems that are selected in the habitat model, but also any other constraints in the model that are represented by the ancillary data layers.

Credits: U.S. Geological Survey (USGS) - Gap Analysis Project (GAP), 2018, California Red-legged Frog (*Rana draytonii*) aCRLFx\_CONUS\_2001v1 Habitat Map: U.S. Geological Survey data release, <https://doi.org/10.5066/F7T43RCM>.

**Data Source:** USGS

**File Name:** california\_red\_legged\_frog\_habitat.tif

---

## LOGGERHEAD SHRIKE

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** This layer shows highly suitable habitats for the reproduction and feeding of Loggerhead Shrike (*Lanius ludovicianus*) within the species' range.

**Data Resolution:** 30m Raster

**Data Units:** Binary, 0 (not suitable) and 1 (suitable)

**Creation Method:** CWHR classifications are based on a combination of FVEG canopy cover, size class and vegetation data. The vegetation data includes a variety of tree, shrub, grassland, and water dominated habitats. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction or feeding of that species in the California Wildlife Habitat Relationship database.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** Loggerhead\_Shrike\_suitable\_habitat.tif

---

## NUTTALL'S WOODPECKER

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** This layer shows highly suitable habitats for the reproduction and feeding of Nuttall's Woodpecker (*Dryobates nuttallii*) within the species' range.

**Data Resolution:** 30m Raster

**Data Units:** Binary, 0 (not suitable) and 1 (suitable)

**Creation Method:** CWHR classifications are based on a combination of FVEG canopy cover, size class and vegetation data. The vegetation data includes a variety of tree, shrub, grassland, and water dominated habitats. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction or feeding of that species in the California Wildlife Habitat Relationship database.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** Nuttalls\_Woodpecker\_suitable\_habitat.tif

---

RINGTAIL CAT

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** This layer shows highly suitable habitats for the reproduction and feeding of Ringtail Cat (*Bassariscus astutus*) within the species' range.

**Data Resolution:** 30m Raster

**Data Units:** Binary, 0 (not suitable) and 1 (suitable)

**Creation Method:** CWHR classifications are based on a combination of FVEG canopy cover, size class and vegetation data. The vegetation data includes a variety of tree, shrub, grassland, and water dominated habitats. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction or feeding of that species in the California Wildlife Habitat Relationship database.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** Ringtail\_Cat\_suitable\_habitat.tif

---

LEAST BELL'S VIREO

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** This layer shows highly suitable habitats for the reproduction and feeding of Least Bell's Vireo (*Vireo bellii pusillus*) within the species' range.

**Data Resolution:** 30m Raster

**Data Units:** Binary, 0 (not suitable) and 1 (suitable)



**Creation Method:** CWHR classifications are based on a combination of FVEG canopy cover, size class and vegetation data. The vegetation data includes a variety of tree, shrub, grassland, and water dominated habitats. Species are considered present, and habitats considered suitable for each 30m cell for which the canopy cover-size-vegetation combination have been deemed highly suitable for the reproduction or feeding of that species in the California Wildlife Habitat Relationship database.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** Least\_Bells\_Vireo\_suitable\_habitat.tif

## COMMUNITY INTEGRITY

The ability of communities to adapt to changing ecological, social, and economic conditions. This entails the capability of an ecological system to sustain a community of organisms that retains the pre-settlement species composition, diversity, and functional organization of natural habitats within a region.

---

## FUNCTIONAL GROUP SPECIES RICHNESS

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** Functional groups are sets of species that share life history characteristics that perform particular functions within an ecosystem. The six functional groups are represented and include a range of trophic levels and ecosystem services. A primary consideration in management is to maintain conditions, adapt to changing conditions and transition to alternate but still productive conditions over time. The maintenance of ecosystem services is a primary concern with climate change.

**Data Resolution:** 30m Raster

**Data Units:** Number of species

**Creation Method:** Species list created from CWHR is divided into six functional groups based on literature. The six functional groups include herbivores, predators, insectivores, soil aerators, seed/spore dispersers and cavity nesters/excavators. The diversity of each functional group is first determined by the number of species for which a given location provides high suitability reproductive habitat (as per species richness calculations). Target conditions can be generated based on percentiles of functional group richness across all patches, so that the 90th percentile or higher is considered in target conditions and the 10th percentile or below is considered to be in a fully departed condition.

**Data Source:**

FVEG

California Department of Fish and Wildlife CWHR version 9.0 (CDFW); 2014

**File Name:** cavity\_nesters\_excavators\_species\_richness.tif; herbivores\_species\_richness.tif;

insectivores\_species\_richness.tif; predators\_species\_richness.tif; seed\_spore\_dispersers\_species\_richness.tif; soil\_aerators\_species\_richness.tif

---

## HABITAT CONNECTIVITY

**Tier:** 1

**Data Vintage:** last updated 08/21/2019

**Metric Definition and Relevance:** The Terrestrial Connectivity dataset is one of the four key components of the California Department of Fish and Wildlife's (CDFW) Areas of Conservation Emphasis (ACE) suite of terrestrial conservation information. The dataset summarizes the relative ability of a species to move across the landscape between patches of suitable habitat. It shows a compilation of linkages, corridors, and natural landscape blocks identified in statewide and regional connectivity studies. Each hexagon (2.5 mi<sup>2</sup>) is ranked into one of the following categories based on the identification of corridors and linkages in statewide, regional, and species-movement studies:

- **5: Irreplicable and Essential Corridors** – The Nature Conservancy's (TNC) Omniscape model identifies channelized areas and priority species movement corridors. The mapped channelized areas are those areas where surrounding land use and barriers are expected to funnel, or concentrate, animal movement. These areas may represent the last available connection(s) between two areas, making them high priority for conservation.
- **4: Conservation Planning Linkages** – Habitat connectivity linkages are often based on species-specific models and represent the best connections between core natural areas to maintain habitat connectivity. Linkages have more implementation flexibility than irreplaceable and essential corridors; any linkage areas not included in rank 5 are included here.
- **3: Connections with Implementation Flexibility** – Areas identified as having connectivity importance but not identified as channelized areas, species corridors or habitat linkage at this time. Future changes in surrounding land use or regional specific information may alter the connectivity rank. Included in this category are areas mapped in the TNC Omniscape study as 'intensified', core habitat areas, and areas on the periphery of mapped habitat linkages.
- **2: Large Natural Habitat Areas** – Large blocks of natural habitat (> 2000 acres) where connectivity is generally intact. This includes natural landscape blocks from the 2010 CEHC and updated with the 2016 Statewide Intactness dataset. Areas mapped as CEHC NLB and not include in the previous ranks, are included here.
- **1: Limited Connectivity Opportunity** – Areas where land use may limit options for providing connectivity (e.g., agriculture, urban) or no connectivity importance has been identified in models. Includes lakes. Some DOD lands are also in this category because they have been excluded from models due to lack of conservation opportunity, although they may provide important connectivity habitat.

**Data Resolution:** 30m Raster

**Data Units:** Categorical; 5 (listed above)

**Creation Method:** Developed by CDFW, the Terrestrial Connectivity dataset summarizes information on terrestrial connectivity by ACE hexagon (2.5 mi<sup>2</sup>) including the presence of mapped corridors or linkages and the juxtaposition to large, contiguous, natural areas. This dataset was developed to support conservation planning efforts by allowing the user to spatially evaluate the relative contribution of an area to terrestrial connectivity based on the results of statewide, regional, and other connectivity analyses. This map builds on the 2010 California Essential

Habitat Connectivity (CEHC) map, based on guidance given in the 2010 CEHC report. The data are summarized by ACE hexagon.

The ACE Terrestrial Connectivity polygon has been converted to 30m Raster and the connectivity description attribute (HabDesc) is classified into the five connectivity ranks (detailed above).

**Data Source:** California Department of Fish and Wildlife; Terrestrial Connectivity, Areas of Conservation Emphasis (ACE), version 3.1

**File Name:** HabitatConnectivity\_2019.tif

## ECONOMIC DIVERSITY

Economic Diversity increases business opportunities that provide regional economic vitality and additional benefits to rural and vulnerable populations. Ecosystem services and forest products provide a foundation for many local and regional economic activities and employment opportunities. Forest management should support a sustainable natural resource-based economy.

**DESIRED OUTCOME:** Forest management and outdoor activities support a sustainable, natural-resource-based economy, particularly in rural communities.

## WOOD PRODUCT INDUSTRY

The wood product industry, with some exceptions (e.g. Big Creek Lumber in Davenport, Pacific Coast Lumber in Paso Robles), is largely absent from the Central Coast Region. However, restoration activities, including vegetation management, are necessary and require financial investments to make progress. This work brings jobs and income to local communities.

---

## COST OF POTENTIAL TREATMENTS

**Tier:** 2

**Data Vintage:** 2023

**Metric Definition and Relevance:** The principle method for maintaining or restoring resilience to the Central Coast landscape involves vegetation treatments. There are many variations on treatments involving different kinds of equipment and different activities of managing vegetation. The metric has gathered available information on the costs of the major treatment methods and incorporated this information into a geospatial database.

There are no treatments of vegetation in the Central Coast that generate revenue. All treatments included here are represented simply as costs per acre.

**Field definitions:**

Mastication = CALFIRE estimates for treatments per acre (Brush = \$1,669, Herbaceous = \$1,813, Woodland = \$1,198, Forest = \$1,788)

Masticat\_1 = USFS estimates per acre (low end = \$800), depends on amount of vegetation

Masticat\_2 = USFS estimates per acre (high end = \$1700), depends on amount of vegetation

Thinning\_m = CALFIRE estimates for manual thinning per acre (Brush = \$2,534, Herbaceous = \$1,851, Woodland = \$2,683, Forest = \$1,461)

Thinning\_1 = USFS estimates per acre (low end = \$450), depends on amount of vegetation

Thinning\_2 = USFS estimates per acre (high end = \$950), depends on amount of vegetation

Thinning\_3 = CALFIRE estimates mechanical thinning per acre (Brush = \$2,500, Herbaceous = N/A, Woodland = \$2,807, Forest = \$957)

Thinning\_4 = USFS estimates mechanical thinning per acre (low end = \$945), depends on amount of vegetation

Thinning\_5 = USFS estimates mechanical thinning per acre (high end = \$1,800), depends on amount of vegetation

Piling\_man = CALFIRE estimates manual piling per acre (Brush = \$2,551, Herbaceous = N/A, Woodland = N/A, Forest = \$1,071)

Piling\_m\_1 = USFS estimates manual piling per acre (low end = \$400), depends on amount of vegetation

Piling\_m\_2 = USFS estimates manual piling per acre (high end = \$1,200), depends on amount of vegetation

Piling\_mec = CALFIRE estimates mechanical piling per acre (Brush = \$1,521, Herbaceous = N/A, Woodland = \$251, Forest = \$640)

Piling\_m\_3 = USFS estimates mechanical piling per acre (low end = \$800), depends on amount of vegetation

Piling\_m\_4 = USFS estimates mechanical piling per acre (high end = \$1,200), depends on amount of vegetation

LopScatter = CALFIRE estimates lop and scatter per acre (Brush = \$1,263, Herbaceous = N/A, Woodland = \$1,217, Forest = \$1,616)

LopScatt\_1 = USFS estimates lop and scatter per acre N/A

LopScatt\_2 = USFS estimates lop and scatter per acre N/A

Herbicide\_ = CALFIRE estimates herbicide (post-treatment) per acre (Brush = \$675, Herbaceous = \$396, Woodland = \$667, Forest = \$325)

Herbicide1 = USFS estimates herbicide (post-treatment) per acre (low end = \$250), depends on amount of vegetation

Herbicid\_1 = USFS estimates herbicide (post-treatment) per acre (high end = \$450), depends on amount of vegetation

Pileburn\_C = CALFIRE estimates pile burn per acre (Brush = \$2,303, Herbaceous = \$3,125, Woodland = N/A, Forest = \$810)

Pileburn\_U = USFS estimates lop and scatter per acre N/A

Pileburn\_1 = USFS estimates lop and scatter per acre N/A

**Data Resolution:** 30m Raster

**Data Units:** Dollars per acre

**Creation Method:** Multiple land managers (Forest Service, CALFIRE) were contacted to obtain current estimates of costs of different treatment methods. We received current estimates from both on treatment costs per acre for a variety of treatment methods. Those cost estimates varied by vegetation type and treatment method. These data were linked to the updated FVEG spatial data and rolled up into a single raster with attributes reflecting these two

cost variables. These data are subject to further refinement and changes in costs. Data will continue to be gathered to improve these estimates.

**Data Source:**

CALFIRE

USDA Forest Service

**File Name:** cost\_per\_acre\_vegtype.tif

## CARBON SEQUESTRATION

Forests play an important role in mitigating climate by sequestering and storing large amounts of carbon. However, forests are at risk of losing carbon because of rates of decay and disturbance, especially with high severity wildfires. Knowing where carbon exists provides a context for where changes in forest conditions will have the greatest impact on carbon storage and sequestration objectives.

**DESIRED OUTCOME:** Carbon sequestration is enhanced in a stable and sustainable manner that yields multiple ecological and social benefits.

Note that all values for carbon have been expressed in Mg C/ha, the international standard for how carbon is measured. If needed, to convert back to the native short tons per acre, divide the Mg/ha by 2.2417023114334.

### CARBON STORAGE

Carbon storage in forest biomass is an essential attribute of stable forest ecosystems and a key link in the global carbon cycle. After carbon dioxide is converted into organic matter by photosynthesis, carbon is stored in forests for a period of time before it is ultimately returned to the atmosphere through respiration and decomposition or disturbance (e.g., fire). A substantial pool of carbon is stored in woody biomass (roots, trunks, branches). Another portion eventually ends up as organic matter in forest floor litter and in soils. Soil carbon does not change very quickly and is difficult to measure directly.

---

### TOTAL ABOVEGROUND CARBON

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Identifying ecosystem carbon is essential to land managers and the Total Aboveground Carbon metric provides an estimate of the amount of existing carbon and its location on California's landscape. The metric also serves to provide context for the other metrics used to quantify carbon sequestration. For example, instability or lack of resilience in forests with low total aboveground carbon would be of less concern than the same degree of instability in a forest that has large total aboveground carbon.

**Data Resolution:** 30m Raster

**Data Units:** Grams dry matter/m<sup>2</sup>

**Creation Method:** The Center for Ecosystem Climate Solutions (CECS) DataEngine model tracks monthly carbon in multiple pools from 1986 to 2021. The carbon components are initialized with eMapR (see [Additional Resources](#)) observations for the early Landsat era; the model then runs freely based on Landsat and other observations. Disturbances and disturbance intensity are tracked annually by Landsat (see other metrics developed by CECS) and

used to quantitatively transfer or combust pools. The model allocates and turns over material based on allometry scaling theory, as adjusted by observational data sets. Aboveground pools (live tree, live shrubs and dead material) are summed for September of 2021.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** CStocks\_Total\_Above\_2021.tif

## CARBON STABILITY

Carbon stability is an important feature in carbon sequestration calculations because carbon turnover – high levels of loss, even if followed by high rates of sequestration – are not as ecologically beneficial as high residency rates for carbon and larger pool values, particularly when stored in large live trees which have many other ecological benefits. The carbon in dead biomass is considered a more unstable component of the carbon pool itself, and a potential destabilizing factor for the live carbon pool in fire-adapted forest ecosystems, especially where it exceeds certain thresholds (e.g., over 46 Mg (total biomass)/ha, Stephens et al., 2022).

---

## ABOVEGROUND CARBON TURNOVER TIME

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** The average lifetime of aboveground live and dead carbon in years. Locations where the lifetime or turnover time is longer have more carbon in more stable pools, such as large trees or large coarse woody debris. Locations where the lifetime or turnover time is shorter have more carbon in labile pools, such as live or dead leaves.

**Data Resolution:** 30m Raster

**Data Units:** Years

**Creation Method:** Calculated from the ratio of total aboveground carbon and annual decomposition. Aboveground carbon and annual decomposition are both calculated for 2021 from a Landsat-driven pools and fluxes model, as described for the total aboveground carbon product. Aboveground turnover time does not currently account for carbon losses and removals with combustion or harvest.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** CStocks\_Turnovertime\_2021.tif

## WATER SECURITY

Forests serve as natural water collection, storage, filtration, and delivery systems as water flows from forests into rivers providing critical aquatic and wetland habitat, while also supplying water for drinking and agriculture. From a more mechanistic perspective, the energy and water balance of forest ecosystems are fundamentally linked. Water is essential to photosynthesis and the latent energy exchange of transpiration is a major driver of water loss. In short, the fate of forests directly influences the quantity and quality of California's freshwater supply.

**DESIRED OUTCOME:** Watersheds provide a reliable supply of clean water despite wide swings in annual precipitation, droughts, flooding, and wildfire.

## QUANTITY

Understanding the interaction between water supply and ecosystem demand informs both the extent of moisture stress and the amount of water available for storage.

---

#### ACTUAL EVAPOTRANSPIRATION TO PRECIPITATION FRACTION DURING DROUGHT

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Plants respond to conditions in their immediate vicinity. Thus, to understand the vegetative moisture stress during drought, it is important to measure the local moisture balance. The actual evapotranspiration fraction (AETF) provides such a measure. Specifically, it indicates whether a location is expected to experience local drying during a drought, or whether the location receives sufficient precipitation that it will remain moist even during an extended drought. An extended drought is defined by a 48-month period where the Standardized Precipitation Index (SPI, NCAR 2022) is two standard deviations below the long-term mean (SPI-48 = negative 2). Such a drought is expected approximately once every 50 years in the Central Coast Region. The AETF ranges from 0 to > 1; a low value indicates a wetter location during drought and a high value indicates a drier location. Locations <1 would be expected to generate runoff, even during a significant drought (SPI-48 drought = negative 2.0), and would be expected to continue generating runoff. Locations > 1 would be expected to desiccate the soil during drought, with negligible runoff, and increasing vegetation drought stress. AET/P does not account for lateral water inflow, either as runoff or irrigation.

**Data Resolution:** 30m Raster

**Data Units:** Dimensionless fraction (AET in mm/P in mm).

**Creation Method:** Calculated as the ratio of actual evapotranspiration (AET) during 2021 Water Year (WY) and precipitation that would be expected for each pixel under a significant drought (SPI-48 drought = negative 2.0). AET is calculated based on Landsat observations and eddy covariance, along with information on local monthly irradiance that accounts for Top of Atmosphere and topographic effects. The AET calculated for 2021 is then divided by the precipitation that would be expected for each pixel under a significant drought (SPI-48 drought = negative 2.0). This quantity of precipitation is calculated for each pixel based on local, down-scaled PRISM data for 1991-2020. This fraction provides a measure of the local water balance during drought, with the higher values indicating a drier location. See <https://doi.org/10.1029/2012JG002027> and <https://doi.org/10.1073/pnas.1319316111> for further information.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** WaterFlux\_AETFrac\_SPI-2\_2021.tif

---

#### PRECIPITATION MINUS ACTUAL EVAPOTRANSPIRATION DURING AVERAGE CONDITIONS

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Runoff is a measure of the water available for storage. It is determined by both the water supply and the demand of the existing vegetation. Annual mean runoff measures the “average” vegetative demand and thus provides a comparative index on the potential available runoff. Specifically, Annual Mean Runoff is the expected surplus water that would discharge to surface or groundwater flows during a series of years with average precipitation. Larger values indicate more runoff under mean conditions.

**Data Resolution:** 30m Raster

**Data Units:** mm/y

**Creation Method:** The Center for Ecosystem Climate Solutions at UC Irvine (CECS) is working with the State and Federal governments in developing scientifically rigorous, stakeholder-informed methods that have produced tailored, integrated data for land management decision makers. The CECS DataEngine model tracks monthly water balance from 1986 to 2021. The Annual Mean Runoff layer is calculated using this CECS DataEngine model logic forced with a series of 4 years that each received precipitation according to the timing and magnitude of the 30-year climate Normal Precipitation (SPI = 0 by definition).

The model water inputs are determined from downscaled PRISM gridded datasets (<https://prism.oregonstate.edu/>). In the case of the Annual Mean Runoff, this reflects the monthly 30 year Normal for each pixel calculated for 1991-2020. Actual evapotranspiration (AET) is calculated from Landsat observations and eddy covariance during 2021, along with information on local monthly irradiance that accounts for Top of Atmosphere (TOA) and topographic effects, as well as monthly temperature and drought stress. Precipitation Minus Actual Evapotranspiration is calculated as the difference; it provides an excellent measure of the long-term runoff from upland pixels. Areas with a higher P-ET produce greater runoff, and areas with a low P-ET tend to produce little or no runoff during average or dry years. See <https://doi.org/10.1029/2012JG002027> and <https://doi.org/10.1073/pnas.1319316111> for further information.

**Data Source:** CECS; <https://california-ecosystem-climate.solutions/>

**File Name:** WaterFlux\_Runoff\_SPI0\_2021.tif

## QUALITY

Understanding the interaction between water supply and ecosystem demand informs both the extent of moisture stress and the amount of water available for storage.

---

## PERCENT IMPERVIOUS SURFACE

**Tier:** 1

**Data Vintage:** 2019

**Metric Definition and Relevance:** This National Land Cover Database (NLCD) product represents urban impervious surfaces as a percentage of developed surface over every 30-meter pixel of California, extracted from a nationwide layer. The definition of impervious means water does not seep into the ground, it runs off into storm sewers and then into local creeks. Examples of impervious surfaces include highways, streets and pavement, driveways, and house roofs. The relevance of impervious surfaces is the higher the proportion of impervious surfaces the more likely flooding can occur.

**Data Resolution:** 30m Raster

**Data Units:** Percent Imperviousness

**Creation Method:** The NLCD 2019 design aims to provide consistent and robust methodologies for production of a multi-temporal land cover and land cover change database from 2001 to 2019 at 2–3-year intervals. Comprehensive research was conducted and resulted in developed strategies for NLCD 2019: continued integration between impervious surface and all landcover products with impervious surface being directly mapped



as developed classes in the landcover, a streamlined compositing process for assembling and preprocessing based on Landsat imagery and geospatial ancillary datasets; a multi-source integrated training data development and decision-tree based land cover classifications; a temporally, spectrally, and spatially integrated land cover change analysis strategy; a hierarchical theme-based post-classification and integration protocol for generating land cover and change products; a continuous fields biophysical parameters modeling method; and an automated scripted operational system for the NLCD 2019 production. For information see [Data | Multi-Resolution Land Characteristics \(MRLC\) Consortium](#)

**Data Source:** National Land Cover Database (NLCD)

**File Name:** nlcd\_2019\_imperviousPercent\_CA.tif

## AIR QUALITY

The goal of healthier forests is aligned with the goal of having healthier air (Cisneros et al., 2014, Long et al., 2018). Forests with sustainable fuel loads create less emissions overall, and support less rapid fire growth, which reduces emissions per day and decreases the chances that smoke from a wildland fire event will create long duration, intense smoke episodes like those we've seen at regional scales during the past decade. Key to supporting the proactive management of smoke and minimization of impacts is a granular understanding at the project scale of where the fuels are, and what potential emissions might occur under wildfire and/or Rx fire scenarios. Those emissions (e.g., from maps like those produced by F3 below) combined with estimates of daily spread can be used to inform operational or scenario-based dispersion modeling (and would be compatible with California's PFIRS smoke management system), which in turn would help fire and air managers better understand where smoke is likely to go, and help inform the public where and when it's likely to occur at potentially unhealthy concentrations.

Tradeoffs between wildfire and Rx fire smoke production (daily, or in total) could be quantified on a first order basis by summing daily or total emissions from high severity vs moderate severity over the area of the respective fire spread polygons. Note that Rx fire smoke impacts are not only different due to per acre differences in emissions, but because the per day emissions can also differ quite substantially. Those emissions numbers could also inform dispersion modeling scenarios showing the relative differences in smoke impacts between wildfire and prescribed scenarios, or even between different wildfire management scenarios.

**DESIRED OUTCOME:** Emissions from fires are limited to primarily low- and moderate-severity fires in wildland ecosystems. Forests improve air quality by capturing pollutants.

## PARTICULATE MATTER

Particle pollution represents a main component of wildfire smoke and the principal public health threat. Fine particles (also known as PM<sub>2.5</sub>) are particles generally 2.5 μm in diameter or smaller and represent a main pollutant emitted from wildfire smoke. Fine particles from wildfire smoke are of greatest health concern.

---

### TOTAL POTENTIAL SMOKE EMISSION - HIGH SEVERITY

**Tier:** 1

**Data Vintage:** 2022

**Metric Definition and Relevance:** SpatialFOFEM/LANDFIRE potential smoke emissions reflect the total PM<sub>2.5</sub> that would be expected to be produced for each pixel for a fire burning under either high or low fire severity conditions. High severity conditions are thought to reflect the emissions that would be expected for a severe

summer wildfire and moderate severity conditions are thought to reflect the emissions that would be expected for a prescribed burn. Potential smore emissions do not consider the probability of a fire or the transport of smoke to more distant locations; they only reflect what would happen locally if a pixel were to burn.

**Data Resolution:** 30m Raster

**Data Units:** Pounds of PM2.5 per acre summed over the both flaming and smoldering phases.

**Creation Method:** Calculated with SpatialFOFEM (First Order Fire Effects Model), embedded in FlamMap 6.2. Fuels are LCP and FCCS 2022 from LANDFIRE (LCP\_LF2022\_FBFM40\_220\_CONUS and LF2022\_FCCS\_220\_CONUS). Conditions for high severity are Pacific coast, Summer, 75% branch consumed, Fuel moisture 20% Duff, 6% 10 hr, 10% 1000 hr, based on very dry conditions per FOFEM 6-7 user guide, page 79 and also FOFEM6 tutorial Sierra burn example ([https://www.firelab.org/sites/default/files/2021-02/FOFEM6\\_Tutorial.pdf](https://www.firelab.org/sites/default/files/2021-02/FOFEM6_Tutorial.pdf)).

**Data Source:**

SpatialFOFEM data outputs (CECS)

Rocky Mountain Research Station

<https://www.firelab.org/project/fofem-fire-effects-model>

**File Name:** Vulner\_PM25\_Severe\_2022.tif

---

## WETLAND INTEGRITY

Wetlands provide critical habitat, store carbon, enhance water quality, control erosion, filter and retain nutrient pollution, and provide spaces for recreation. They are local and regional centers of biodiversity, and support species found nowhere else across western landscapes. Functional wetland ecosystems will serve increasingly important roles in buffering impacts from extreme climate events, and upland disturbances such as flooding and erosion. Meadow and riparian ecosystems provide ecosystem services and are key linkages between upland and aquatic systems in forested landscapes.

**DESIRED OUTCOME:** Wetland ecosystems are biologically intact, provide multiple ecosystem services, and meadow and riparian ecosystems provide key linkages between upland and aquatic systems in forested landscapes.

## COMPOSITION

Wetland composition pertains to the array of different wetland types, their relative abundance, the uniqueness of their co-occurrence and composition, and their integrity in a given location and area within and across landscapes. Wetland ecosystems include all lentic (e.g. lakes, ponds, bogs, fens) and lotic (e.g., rivers, streams, springs, seeps) aquatic ecosystems, as well as associated vegetated wetlands such as wet meadows and riparian vegetation.

---

## AQUATIC SPECIES RICHNESS

**Tier:** 1

**Data Vintage:** 2018

**Metric Definition and Relevance:** Aquatic native species richness is a measure of species biodiversity, and is one measurement used to describe the distribution of overall species biodiversity in California for the California Department of Fish and Wildlife (CDFW) Areas of Conservation Emphasis Project (ACE). Native species richness represents a count of the total number of native aquatic species potentially present in each watershed based on species range and distribution information. The data can be used to view patterns of species diversity, and to identify areas of highest native richness across the state. The species count consists of four taxonomic groups – fish, aquatic invertebrates, aquatic amphibians, and aquatic reptiles.

**Data Resolution:** 30m Raster

**Data Units:** Count

**Creation Method:** For more information, see the Aquatic Native Species Richness Factsheet (2018) at <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=150852>

The California Department of Fish and Wildlife (CDFW) Areas of Conservation Emphasis (ACE) is a compilation and analysis of the best-available statewide spatial information in California on biodiversity, rarity and endemism, harvested species, significant habitats, connectivity and wildlife movement, climate vulnerability, climate refugia, and other relevant data (e.g., other conservation priorities such as those identified in the State Wildlife Action Plan (SWAP), stressors, land ownership). ACE addresses both terrestrial and aquatic data.

**Data Source:**

Aquatic Native Species Richness Summary, Areas of Conservation Emphasis (ACE), version 3.0, California Department of Fish and Wildlife (CDFW)  
ACE database

**File Name:** aquatic\_species\_richness.tif

---

## WETLAND DIVERSITY

**Tier:** 1

**Data Vintage:** 2018

**Metric Definition and Relevance:** This data set represents the extent, approximate location, and type of wetlands and deepwater habitats in California. These data delineate the areal extent of wetlands and surface waters as defined by Cowardin et al. (1979).

**Data Resolution:** 30m raster

**Data Units:** Thematic

**Creation Method:** Downloaded from the National Wetlands Inventory (NWI), polygon converted to 30 meter raster. For more information see <https://www.fws.gov/program/national-wetlands-inventory>.

**Definition of values:**

- Lake = Lake or reservoir basin. Lacustrine wetland and deepwater (L).
- Freshwater Emergent Wetland = Herbaceous marsh, fen, swale and wet meadow. Palustrine emergent (PEM).
- Estuarine and Marine Wetland = Vegetated and non-vegetated brackish and saltwater marsh, shrubs, beach, bar, shoal or flat. Estuarine intertidal and Marine intertidal wetland (E2, M2).

- Other = Farmed wetland, saline seep and other miscellaneous wetland. Palustrine wetland (Misc. types, PUS, Pf..)
- Freshwater Pond = Pond. Palustrine unconsolidated bottom, Palustrine aquatic bed (PUB, PAB).
- Estuarine and Marine Deepwater = Open water estuary, bay, sound, open ocean. Estuarine and Marine subtidal water (E1, M1).
- Riverine = River or stream channel. Riverine wetland and deepwater (R).
- Freshwater Forested/Shrub Wetland = Forested swamp or wetland shrub bog or wetland. Palustrine forested and/or Palustrine shrub (PFO, PSS).

**Data Source:** The National Wetlands Inventory, US Fish & Wildlife Service (USFWS)

**File Name:** NWI\_WetlandsType\_2018\_30m.tif

## RIPARIAN HABITAT

**Tier:** 1

**Data vintage:** 2019

**Metric Definition and Relevance:** These data depict 10-meter raster riparian areas for 50-year flood heights for California in 2019.

**Data Resolution:** 10m Raster

**Data Units:** binary

**Creation Method:** Fifty-year flood heights were estimated using U.S. Geological Survey (USGS) stream gage information. NHDPlus version 2.1 was used as the hydrologic framework to delineate riparian areas. The U.S. Fish and Wildlife Service’s National Wetland Inventory and USGS 10-meter digital elevation models were also used in processing these data. See <https://doi.org/10.2737/RDS-2019-0030>

**Credits:** Sinan Abood, Ph.D. GISP; Research Scientist, Forest Service Washington Office (WO) – Biological & Physical Resources (BPR)

**Data Source:** USDA Forest Service

**File Name:** RiparianAreas10m\_2019.tif

## SOCIAL AND CULTURAL WELL-BEING

The landscape provides a place for people to connect with nature, recreate, to maintain and improve their overall health, and an opportunity to contribute to environmental stewardship. While the elements of this pillar include public health and engagement, recreation quality, and equitable opportunities producing quantifiable, measurable and actionable metrics remains challenging. These metrics are still under development and insights into these potential metrics are appreciated.

**DESIRED OUTCOME:** The landscape provides a place for people to connect with nature, to recreate, to maintain and improve their overall health, and to contribute to environmental stewardship, and is a critical component of their identity.

## ENVIRONMENTAL JUSTICE

Environmental Justice is the fair treatment and meaningful involvement of all people regardless of race, color, national origin or income regarding the development, implementation and enforcement of environmental laws, regulations, policies and land management.

---

## POVERTY PERCENTILE

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Percent of population living below two times the federal poverty level. The U.S. Census Bureau determines the federal poverty level each year. The poverty level is based on the size of the household and the age of family members. If a person or family's total income before taxes is less than the poverty level, the person or family are considered in poverty. Many studies have found that people living in poverty are more likely than others to become ill from pollution.

**Data Resolution:** 30m Raster

**Data Units:** percentile

**Creation Method:** CalEnviroScreen, Version 4.0, is a science-based method for identifying impacted communities by taking into consideration pollution exposure and its effects, as well as health and socioeconomic status, at the census-tract level. CalEnviroScreen 4.0 uses the census tract as the unit of analysis. Census tract boundaries are available from the Census Bureau. CalEnviroScreen uses the Bureau's 2010 boundaries. New boundaries will be drawn by the Census Bureau as part of the 2020 Census but will not be available until after 2022. OEHHA will address updates to census tract geography in CalEnviroScreen at that time. There are approximately 8,000 census tracts in California, representing a relatively fine scale of analysis. Census tracts are made up of multiple census blocks, which are the smallest geographic unit for which population data are available. Some census blocks have no people residing in them (unpopulated blocks).

The poverty percentile is derived from

- The 2015-2019 American Community Survey, a dataset containing the number of individuals below 200 percent of the federal poverty level was downloaded by census tracts for the state of California.
- The number of individuals below 200% of the poverty level was divided by the total population for whom poverty status was determined.
- Unlike the US Census, ACS estimates come from a sample of the population and may be unreliable if they are based on a small sample or population size. The standard error (SE) and relative standard error (RSE) were used to evaluate the reliability of each estimate.
- The SE was calculated for each census tract using the formula for approximating the SE of proportions provided by the ACS (American Community Survey Office, 2013, pg. 13, equation 4). CalEnviroScreen 4.0 189 When this approximation could not be used, the formula for approximating the SE of ratios (equation 3) was used instead.
- The RSE is calculated by dividing a tract's SE by its estimate of the percentage of the population living below twice the federal poverty level, and taking the absolute value of the result.
- Census tract estimates that met either of the following criteria were considered reliable and included in the analysis:
  - RSE less than 50 (meaning the SE was less than half of the estimate) OR
  - SE was less than the mean SE of all California census tract estimates for poverty.

- Census tracts with unreliable estimates received no score for the indicator (null). The indicator was not factored into that tract's overall CalEnviroScreen score.
- Census tracts that met the inclusion criteria were sorted and assigned percentiles based on their position in the distribution.

**Data Source:** California Environmental Protection Agency, CalEnviroScreen 4.0

**File Name:** Poverty\_Pctl.tif

## HOUSING BURDEN PERCENTILE

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Housing-Burdened Low-Income Households. Percent of households in a census tract that are both low income (making less than 80% of the HUD Area Median Family Income) and severely burdened by housing costs (paying greater than 50% of their income to housing costs). (5-year estimates, 2013-2017).

The cost and availability of housing is an important determinant of well-being. Households with lower incomes may spend a larger proportion of their income on housing. The inability of households to afford necessary non-housing goods after paying for shelter is known as housing-induced poverty. California has very high housing costs relative to much of the country, making it difficult for many to afford adequate housing. Within California, the cost of living varies significantly and is largely dependent on housing cost, availability, and demand.

Areas where low-income households may be stressed by high housing costs can be identified through the Housing and Urban Development (HUD) Comprehensive Housing Affordability Strategy (CHAS) data. We measure households earning less than 80% of HUD Area Median Family Income by county and paying greater than 50% of their income to housing costs. The indicator takes into account the regional cost of living for both homeowners and renters, and factors in the cost of utilities. CHAS data are calculated from US Census Bureau's American Community Survey (ACS).

**Data Resolution:** 30m Raster

**Data Units:** Percent

**Creation Method:** CalEnviroScreen, Version 4.0, is a science-based method for identifying impacted communities by taking into consideration pollution exposure and its effects, as well as health and socioeconomic status, at the census-tract level. CalEnviroScreen 4.0 uses the census tract as the unit of analysis. Census tract boundaries are available from the Census Bureau. CalEnviroScreen uses the Bureau's 2010 boundaries. New boundaries will be drawn by the Census Bureau as part of the 2020 Census but will not be available until 2022. OEHHA will address updates to census tract geography in CalEnviroScreen at that time. There are approximately 8,000 census tracts in California, representing a relatively fine scale of analysis. Census tracts are made up of multiple census blocks, which are the smallest geographic unit for which population data are available. Some census blocks have no people residing in them (unpopulated blocks).

The CalEnviroScreen model is based on the CalEPA working definition in that:

- The model is place-based and provides information for the entire State of California on a geographic basis. The geographic scale selected is intended to be useful for a wide range of decisions.

- The model is made up of multiple components cited in the above definition as contributors to cumulative impacts.
- The model includes two components representing Pollution Burden – Exposures and Environmental Effects
- The model includes two components representing Population Characteristics – Sensitive Populations (e.g., in terms of health status and age) and Socioeconomic Factors.

The American Community Survey (ACS) is an ongoing survey of the US population conducted by the US Census Bureau and has replaced the long form of the decennial census. Unlike the decennial census, which attempts to survey the entire population and collects a limited amount of information, the ACS releases results annually based on a sub-sample of the population and includes more detailed information on socioeconomic factors. Multiple years of data are pooled together to provide more reliable estimates for geographic areas with small population sizes. Each year, the HUD receives custom tabulations of ACS data from the US Census Bureau. These data, known as the "CHAS" data (Comprehensive Housing Affordability Strategy), demonstrate the extent of housing problems and housing needs, particularly for low-income households. The most recent results available at the census tract scale are the 5-year estimates for 2013-2017. The data are available from the HUD user website (see page 174 in the document link below:

<https://oehha.ca.gov/media/downloads/calenviroscreen/report/calenviroscreen40reportf2021.pdf>

**Data Source:** California Environmental Protection Agency, CalEnviroScreen 4.0

**File Name:** HousingBurdenPctl.tif

## UNEMPLOYMENT PERCENTILE

**Tier:** 1

**Data Vintage:** 2021

**Metric Definition and Relevance:** Percentage of the population over the age of 16 that is unemployed and eligible for the labor force. Excludes retirees, students, homemakers, institutionalized persons except prisoners, those not looking for work, and military personnel on active duty (5-year estimate, 2015-2019).

Because low socioeconomic status often goes hand-in-hand with high unemployment, the rate of unemployment is a factor commonly used in describing disadvantaged communities. On an individual level, unemployment is a source of stress, which is implicated in poor health reported by residents of such communities. Lack of employment and resulting low income often constrain people to live in neighborhoods with higher levels of pollution and environmental degradation.

**Data Resolution:** 30m Raster

**Data Units:** Percent

**Creation Method:** CalEnviroScreen, Version 4.0, is a science-based method for identifying impacted communities by taking into consideration pollution exposure and its effects, as well as health and socioeconomic status, at the census-tract level. CalEnviroScreen 4.0 uses the census tract as the unit of analysis. Census tract boundaries are available from the Census Bureau. CalEnviroScreen uses the Bureau's 2010 boundaries. New boundaries will be drawn by the Census Bureau as part of the 2020 Census but will not be available until 2022. OEHHA will address updates to census tract geography in CalEnviroScreen at that time. There are approximately 8,000 census tracts in California, representing a relatively fine scale of analysis. Census tracts are made up of multiple census blocks,

which are the smallest geographic unit for which population data are available. Some census blocks have no people residing in them (unpopulated blocks).

The CalEnviroScreen model is based on the CalEPA working definition in that:

- The model is place-based and provides information for the entire State of California on a geographic basis. The geographic scale selected is intended to be useful for a wide range of decisions.
- The model is made up of multiple components cited in the above definition as contributors to cumulative impacts.
- The model includes two components representing Pollution Burden – Exposures and Environmental Effects
- The model includes two components representing Population Characteristics – Sensitive Populations (e.g., in terms of health status and age) and Socioeconomic Factors.

The American Community Survey (ACS) is an ongoing survey of the US population conducted by the US Census Bureau. Unlike the decennial census, which attempts to survey the entire population and collects a limited amount of information, the ACS releases results annually based on a sub-sample of the population and includes more detailed information on socioeconomic factors such as unemployment. Multiple years of data are pooled together to provide more reliable estimates for geographic areas with small population sizes. The most recent results available at the census tract level are the 5-year estimates for 2015-2019. The data are made available using the U.S. Census data download website.

**Data Source:** California Environmental Protection Agency, CalEnviroScreen 4.0

**File Name:** Unemployment\_Pctl.tif

## OPERATIONAL DATA LAYERS

In addition to the metric data layers assembled for this RRK project, a set of “operational” GIS data layers have been assembled to support use of the metrics. These data provide land use context (e.g. ownership, land use designations, background ecological information (e.g. climate refugia, stream locations, climate classes), infrastructure (roads, operational constraints, powerline corridors), and Forest Service policy information (spotted owl PACs, critical habitat maps for listed species, wilderness/roadless/wild and scenic rivers). These data are provided to assist managers in putting proposed treatments into context for what is feasible and what might constrain project planning.

Data layers provided within this designation of operational data are in their native projection and format with any embedded metadata maintained.

### FIRE

---

#### RECENT FIRE SEVERITY CLASS

**Data Vintage:** 2021



**Metric Definition and Relevance:** Fire severity classification (low, moderate, high) that burned within the last 10 years (2012-2021).

**Data Resolution:** 30m raster

**Data Units:** Value, 1 to 3

**Creation Method:** The difference-adjusted relativized difference normalized burn ratio (RdNBR) was calculated using methods modified from Parks et al (2018). Fire perimeters were obtained from CAL FIRE's April 2021 fire perimeter database. A function for estimating basal area loss from RdNBR values was fit to data from Miller et al (2009) using quasibinomial logistic regression and applied to the 2012-2021 fires. Estimated basal area loss was thresholded to represent low (< 25% loss), moderate (25% – 75% loss), and high (> 75% loss) burn severity. For areas where multiple sequential fires burned from 2012-2021 the maximum burn severity is reported. Updated April 2023 to incorporate CAL FIRE's October-2022 revisions to fire perimeters and to minimize data loss resulting from spatial reprojection.

- 1: Low Severity
- 2: Moderate Severity
- 3: High Severity

**Data Source:**

- Landsat 8, NASA
- Fire History (April 2022), CAL FIRE
- Postfire mortality data, Miller et al. 2009

**File Name:** fire\_severity\_class\_max\_2012to2021\_v2.tif

---

## HOUSING UNIT DENSITY

**Data Vintage:** 2020

**Metric Definition and Relevance:** HUDen is a raster of housing-unit density measured in housing units per square kilometer. The HUDen raster was generated using population and housing-unit count and data from the U.S. Census Bureau, building footprint data from Microsoft, and land cover data from LANDFIRE.

**Data Resolution:** 30m Raster

**Data Units:** Housing units per square kilometer

**Creation Method:** Generate the HUDen raster from the building points. We first converted the building points to a 30-m raster where the raster value is the sum of the housing-units-per-centroid attribute of all building centroids within each raster grid cell. We then generated a smoothed density raster using a three-step process: 1) calculate a 200-m radius moving-window sum of the 30-m housing-unit count raster; 2) calculate a 200-m radius moving-window sum of habitable land cover (in sq km), where habitable land cover is all land covers except open water and permanent-snow/ice; and 3) divide the smoothed housing-unit count raster by the smoothed habitable land cover raster to generate housing unit density in housing units/sq km. To produce the final integer version of the

HUDen raster, we set values less than 0.1 HU/sq km to zero, values between 0.1 and 1.5 to a value of 1 HU/sq km, and rounded all other values to the nearest integer.

**Data Source:** Pyrologix, LLC

**File Name:** HUDen\_2020.tif

---

## WILDLAND URBAN INTERFACE

**Data Vintage:** 2020

**Metric Definition and Relevance:** The wildland urban interface (WUI) is the area where urban development is in close proximity to wildland vegetation. WUI data for the conterminous U.S. based on 125 million building locations where buildings intermingle with or abut wildland vegetation according to the Federal Register definitions of the WUI. According to the definitions used for our building-based maps and for the census-based maps, WUI is where building density exceeds 6.17 units/km<sup>2</sup> and where land cover is either (1) at least 50% wildland vegetation (intermix) or (2) under 50% wildland vegetation but within 2.4 km (1.5 miles) of a patch of wildland vegetation at least 5 km<sup>2</sup> in area that contains at least 75% vegetation (interface). The distance selected for the interface definition is based on research from the California Fire Alliance suggesting that this is the average distance firebrands can travel from an active wildfire front (Stewart et al., 2007).

**Data Resolution:** 30m Raster

**Data Units:** Categorical

**Creation Method:** Building point locations were obtained from a Microsoft product released in 2018, updated to 2019-2020 for most of California, which classified building footprints based on high-resolution satellite imagery. Maps were also based on wildland vegetation mapped by the 2016 National Land Cover Dataset (Yang et al., 2018). The mapping algorithm utilized definitions of the WUI from the U.S. Federal Register (USDA & USDI, 2001) and Radeloff et al. (2005). Both classes required a minimum building density of 6.17 buildings per km<sup>2</sup>. This map of intermix and interface WUI was generated using a circular neighborhood size based on radius distance of 100m to determine building density and vegetation cover on a pixel-by-pixel basis (Bar Massada et al., 2013). Source: USGS ScienceBase Data Catalog; <https://www.sciencebase.gov/catalog/item/617bfb43d34ea58c3c70038f>

Values in the raster are defined as:

- 0: Not WUI
- 1: Intermix WUI
- 2: Interface WUI

**Data Source:** WUI, Carlson et al, 2022

**File Name:** MSB\_WUI\_CA\_100m.tif

---

## ADMINISTRATIVE

---

## URBAN-AGRICULTURE LAND USE

**Definition and Relevance:** This dataset covers the urban and agricultural landscape for all forms of urban and agricultural land use in California. It was created using a combination of best available land cover data from multiple sources (see below). These data are used as a mask for selected metrics in the RRK project where inclusion of urban and agricultural cover potentially creates confusion in calculations of the metric.

**Data Vintage:** FMMP – 2018; NLCD – 2020; MS Bldg – multiple dates

**Data Resolution:** Raster, 30m

**Data Units:** Thematic

**Creation Method:**

1. [Farmland Mapping and Monitoring Program \(FMMP\)](#) land-use data from 2018 was converted to 30m raster as the base input, using the values from the Type field of:
  - Farmland of Statewide Importance
  - Unique Farmland
  - Farmland of Local Importance
  - Urban and Built-Up Land
  - Rural Residential Land
  - Confined Animal Agriculture
2. Secondly, to bring more current data in, [LANDFIRE 2020 Existing Vegetation Type \(EVT\)](#) from 2020 was converted to 30m raster, using the values from EVT group name of:
  - Developed-Low Intensity
  - Developed-Medium Intensity
  - Developed-High Intensity
  - Agriculture-Cultivated Crops and Irrigated Agriculture
3. Lastly, [Building Footprints - Bing Maps \(microsoft.com\)](#) polygons were converted to 30m raster and added to the stack to include the most recent urban footprints.

**Data Source:**

Farmland Mapping and Monitoring Program (FMMP)

LANDFIRE: Existing Vegetation Type, U.S. Department of Agriculture and U.S. Department of the Interior

MS Building Footprints

**File Name:** UrbanAgLanduse\_RRK\_2020.tif

---

## BUILDING STRUCTURE DENSITY

**Definition and Relevance:** A raster dataset containing building footprints of California.

**Data Vintage:** The vintage of the footprints depends on the vintage of the underlying imagery. Bing Imagery is a composite of multiple sources with different capture dates.

**Data Resolution:** Raster, 10m

**Data Units:** binary

**Creation Method:** Vector spatial data called US Building Footprints contained in a Microsoft dataset (available at <https://github.com/microsoft/USBuildingFootprints>) downloaded, clipped to California and converted to a 10m raster. For more information visit: [Building Footprints - Bing Maps \(microsoft.com\)](#)

**Data Source:** MS Building Footprints

**File Name:** CA\_bldgFootprints\_10m.tif

---

## HIGH-USE RECREATION AREAS

**Definition and Relevance:** A recreation site is a discrete area on a National Forest that provides recreation opportunities, receives recreational use, and requires a management investment to operate and/or maintain to standard under the direction of an administrative unit in the National Forest System. Recreation sites range in development from relatively undeveloped areas, with little to no improvements (Development Scale 0 and 1), to concentrations of facilities and services evidencing a range of amenities and investment (Development Scale 2 through 5).

Recreation opportunities are point locations of recreational site activities available to visitors and populates the Forest Service websites (<https://www.fs.usda.gov/>), and the interactive visitor map (<https://www.fs.usda.gov/ivm/>).

**Data Resolution:** Points

**Data Units:** Tabular attributes

**Creation Method:** see Metadata

**Data Source:** USFS Enterprise Data Warehouse (EDW)

**File Name:** RECAREAACTIVITIES\_V\_2023.shp

---

## LAND DESIGNATIONS

**Definition and Relevance:** Wilderness, Roadless, Wild and Scenic River

**Data Vintage:** 2022

**Data Resolution:** ArcGIS file geodatabase: Vector, polygon

**Data Units:** Tabular attributes

**Creation Method:** Data layers pulled from the Enterprise Data Warehouse for land designations:

- *Wilderness* – area designated as a National Wilderness in the National Wilderness Preservation System
- *Inventoried Roadless Areas* – the 2001 Roadless Rule establishes prohibitions on road construction, road reconstruction, and timber harvesting on inventoried roadless areas on National Forest System lands by the following classifications:
  - 1B = Inventoried Roadless Areas where road construction and reconstruction is prohibited
  - 1B-1 = Inventoried Roadless Areas that are recommended for wilderness designation in the forest plan and where road construction and reconstruction is prohibited
  - 1C = Inventoried Roadless Areas where road construction and reconstruction is not prohibited
- *Wild and Scenic Rivers* – area designated as a National Wild, Scenic, or Recreational River within the National Wild and Scenic River System. The designations and definitions are:
  - Wild (W) – Those rivers or sections of rivers that are free of impoundments and generally inaccessible except by trail, with watersheds or shorelines essentially primitive and waters unpolluted. These represent vestiges of primitive America.

- Scenic (S) – Those rivers or sections of rivers that are free of impoundments, with shorelines or watersheds still largely primitive and shorelines largely undeveloped, but accessible in places by roads.
- Recreational (R) – Those rivers or sections of rivers that are readily accessible by road or railroad, that may have some development along their shorelines, and that may have undergone some impoundment or diversion in the past.

**Data Source:** USFS Enterprise Data Warehouse (EDW)

**File Name:** Wilderness\_2023.shp; Roadless\_2001.shp; WildScenicRiver\_2023.shp

## OWNERSHIP

**Definition and Relevance:** Ownership is a commonly used base layer used in a wide range of business functions and these data are intended to provide a depiction of the land ownership within the RRK project area.

**Data Vintage:** FS\_BasicOwnership: 2022, ownership: 2022

**Data Resolution:** Vector, polygon

**Data Units:** Tabular attributes

### Creation Method:

- *FS\_BasicOwnership\_2022.shp* – an area depicted as surface ownership parcels dissolved on the same ownership classification administered by the USDA Forest Service (USFS).
- *ownership22\_1* – California Multi-Source Land Ownership, includes lands owned by each federal agency (including USFS), state agency, local government entities, conservation organizations, and special districts. It does not include lands of private ownership.

**Data Source:** USDA Forest Service, CAL FIRE

**File Name:** FS\_BasicOwnership\_2022.shp; ownership22\_1.shp

## ROADS

**Definition and Relevance:** This California statewide dataset was downloaded from [Geofabrik's free download server](#) for California. This server has data extracts from the OpenStreetMap project which are normally updated every day.

**Data Vintage:** 2022

**Data Resolution:** Vector, line

**Data Units:** Tabular attributes

**Creation Method:** To simplify the layer, major roads were exported with the following selection of the attribute “fclass”:

- 5111 = motorway
- 5112 = trunk
- 5113 = primary
- 5114 = secondary

- 5121 = unclassified
- 5122 = residential
- 5123 = living street

**Data Source:** [Open Street Map](#) roads based on Tiger Lines (OSM)

**File Name:** OSM\_majorRoads\_CA\_2022.shp

## TRANSMISSION LINES

**Definition and Relevance:** This electric transmission line California statewide dataset was downloaded from PG&E (Pacific Gas & Electric) and was subsetted to include only lines less than or equal to 115 kV (kilovolts). This subset was chosen from the original dataset for use in planning because it has been determined (via inspections of PG&E database of fires caused by power lines from 2020-2022) that virtually every fire caused by power lines was from a distribution lines less than 115 kv. Most wildfires caused by power lines are from distribution lines less than 44kv. Thus this database provides information on where those power lines are and can be used to compare with locations that have potential for high severity wildfire.

**Data Vintage:** 2023

**Data Resolution:** Vector, line

**Data Units:** Tabular attributes

**Creation Method:** PG&E’s Integration Capacity Analysis (ICA) map is designed to help contractors and developers find information on potential project sites for distributed energy resources (DERs). ICA is a complex modeling study that uses detailed information about the electric distribution system, which includes items such as physical infrastructure, load performance, and existing and queued generators. The analysis simulates the ability of individual distribution line sections to accommodate additional DERs without potentially causing issues that would impact customer reliability and power quality. Potential issues could result in distribution line upgrade requirements that would impact cost and/or timeline for DER interconnections.

We have selected only those lines that are 115 kv or less to include in this data layer.

Transmission lines:

- Carry electricity across the state
- Transport bulk electricity at high voltages ranging from 60 kV-500 kV
- Are usually supported on tall metal towers, but sometimes on wooden poles
- Have different vegetation standards than distribution lines due to the high voltages they carry
- Are managed using the utility industry’s best-management practice of Wire Zone Border Zone
- Require only low-growing vegetation underneath—typically nothing taller than 10 feet at maturity

[https://www.pge.com/en\\_US/safety/yard-safety/powerlines-and-trees/transmission-vs-distribution-power-lines.page](https://www.pge.com/en_US/safety/yard-safety/powerlines-and-trees/transmission-vs-distribution-power-lines.page)

**Data Source:** PG&E

[PG&E Integration Capacity Analysis and Distribution Investment Deferral Framework maps \(pge.com\)](#)

**File Name:** TransmissionLines\_upTo\_115kv.shp

---

## DISTRIBUTION LINES

**Definition and Relevance:** This electric distribution line California statewide dataset was downloaded from PG&E (Pacific Gas & Electric). This 'FeederDetail' dataset carries voltage under the 'Nominal\_Voltage' attribute for the distribution system, all under 44kV. These distribution lines often can cross wildlands and through vegetated areas and are typically the most likely to be related to a wildfire.

This subset was chosen from the original dataset for use in planning because it has been determined (via inspections of PG&E database of fires caused by power lines from 2020-2022) that virtually every fire caused by power lines was from a distribution lines less than 115 kv. Most wildfires caused by power lines are from distribution lines less than 44kv. Thus this database provides information on where those power lines are and can be used to compare with locations that have potential for high severity wildfire.

**Data Vintage:** 2023

**Data Resolution:** Vector, line

**Data Units:** Tabular attributes

**Creation Method:** PG&E's Integration Capacity Analysis (ICA) map is designed to help contractors and developers find information on potential project sites for distributed energy resources (DERs). ICA is a complex modeling study that uses detailed information about the electric distribution system, which includes items such as physical infrastructure, load performance, and existing and queued generators. The analysis simulates the ability of individual distribution line sections to accommodate additional DERs without potentially causing issues that would impact customer reliability and power quality. Potential issues could result in distribution line upgrade requirements that would impact cost and/or timeline for DER interconnections.

Distribution lines:

- Deliver electricity to neighborhoods and communities over a shorter distance than transmission lines
- Are generally supported by wooden poles and not as high as transmission lines
- Are the final stage of electricity delivery to homes and businesses
- Carry lower voltage electricity that is still powerful enough to cause injury or death
- Trees growing near these lines may be managed with directional pruning, but removal is often best.

[https://www.pge.com/en\\_US/safety/yard-safety/powerlines-and-trees/transmission-vs-distribution-power-lines.page](https://www.pge.com/en_US/safety/yard-safety/powerlines-and-trees/transmission-vs-distribution-power-lines.page)

**Data Source:** PG&E

[PG&E Integration Capacity Analysis and Distribution Investment Deferral Framework maps \(pge.com\)](#)

**File Name:** FeederDetail.shp

---

## CWHR VEGETATION

**Definition and Relevance:** Vegetation maps are important for characterizing many important features of a landscape such as wildlife habitat, fuels conditions, forest composition, and carbon. Such data are most useful if they can depict vegetation type, cover, and tree size class. This version was created to capture current conditions as best as possible through a variety of existing and current sources. Cross-walks were used to compile the various sources into the common classification scheme, the California Wildlife Habitat Relationships (CWHR) system. See CWHR for more details on the CWHR system ([California Wildlife Habitat Relationships](#)).

Key field names in this data set (there are others) are defined as follows:

WHRALL - Unique habitat data label. Concatenated from separate habitat attributes WHRtype, WHRsize and WHRdensity.

WHRNUM - Unique number for each Wildlife Habitat Relationship class (WHRtype).

WHRNAME - Unique name for each Wildlife Habitat Relationship class (WHRtype)

WHRTYPE - Unique Wildlife Habitat Relationship (WHR) class code

WHRSIZE - Wildlife Habitat Relationship Size Class (tree types only)

WHRDENSITY - Wildlife Habitat Relationship class (tree types only)

SOURCE\_NAME - General description of where the source data layer used for a given geography

SOURCE\_YEAR - Year of base imagery that source data layer references for a given geography

WHR Codes for Vegetation Types:

### Tree Dominated Habitats

| CWHR Code | Type Description         |
|-----------|--------------------------|
| ASP       | Aspen                    |
| BOP       | Blue Oak-Foothill Pine   |
| BOW       | Blue Oak Woodland        |
| COW       | Coastal Oak Woodland     |
| CPC       | Closed-Cone Pine-Cypress |
| DFR       | Douglas Fir              |
| DRI       | Desert Riparian          |
| EPN       | Eastside Pine            |



|     |                          |
|-----|--------------------------|
| EUC | Eucalyptus               |
| JPN | Jeffrey Pine             |
| JST | Joshua Tree              |
| JUN | Juniper                  |
| KMC | Klamath Mixed Conifer    |
| LPN | Lodgepole Pine           |
| MHC | Montane Hardwood-Conifer |
| MHW | Montane Hardwood         |
| MRI | Montane Riparian         |
| PJN | Pinyon-Juniper           |
| POS | Palm Oasis               |
| PPN | Ponderosa Pine           |
| RDW | Redwood                  |
| RFR | Red fir                  |
| SCN | Subalpine Conifer        |
| SMC | Sierran Mixed Conifer    |
| VOW | Valley Oak Woodland      |
| VRI | Valley Foothill Riparian |
| WFR | White fir                |

**Shrub Dominated Habitats**

| CWHR Code | Type Description   |
|-----------|--------------------|
| ADS       | Alpine Dwarf-Shrub |

|     |                            |
|-----|----------------------------|
| ASC | Alkali Desert Scrub        |
| BBR | Bitterbrush                |
| CRC | Chamise-Redshank Chaparral |
| CSC | Coastal Scrub              |
| DSC | Desert Scrub               |
| DSS | Desert Succulent Shrub     |
| DSW | Desert Wash                |
| LSG | Low Sage                   |
| MCH | Mixed Chaparral            |
| MCP | Montane Chaparral          |
| SGB | Sagebrush                  |

**Herbaceous Dominated Habitats**

| CWHR Code | Type Description        |
|-----------|-------------------------|
| AGS       | Annual Grass            |
| FEW       | Fresh Emergent Wetland  |
| PAS       | Pasture                 |
| PGS       | Perennial Grass         |
| SEW       | Saline Emergent Wetland |
| WTM       | Wet Meadow              |

**Aquatic Habitats**

| CWHR Code | Type Description |
|-----------|------------------|
|-----------|------------------|

|     |            |
|-----|------------|
| EST | Estuarine  |
| LAC | Lacustrine |
| MAR | Marine     |
| RIV | Riverine   |

**Developed Habitats**

| CWHR Code | Type Description              |
|-----------|-------------------------------|
| CRP       | Cropland                      |
| DGR       | Dryland Grain Crops           |
| DOR       | Deciduous Orchard             |
| EOR       | Evergreen Orchard             |
| IGR       | Irrigated Grain Crops         |
| IRF       | Irrigated Row and Field Crops |
| IRH       | Irrigated Hayfield            |
| OVN       | Orchard - Vineyard            |
| RIC       | Rice                          |
| URB       | Urban                         |
| VIN       | Vineyard                      |

**Non-vegetated Habitats**

| CWHR Code | Type Description |
|-----------|------------------|
| BAR       | Barren           |

WHR Codes for Tree Size Classes:

| CWHR Code | CWHR Size Class    | Conifer Crown Diameter  | Hardwood Crown Diameter | DBH           |
|-----------|--------------------|---|-------------------------|---------------|
| 1         | Seedling tree      | n/a   | n/a                     | <1.0"         |
| 2         | Sapling tree       | n/a   | <15.0'                  | 1.0" - 5.9"   |
| 3         | Pole tree          | <12.0'  | 15.0' - 29.9'           | 6.0" - 10.9"  |
| 4         | Small tree         | 12.0' - 23.9'   | 30.0' - 44.9'           | 11.0" - 23.9" |
| 5         | Medium/large tree  | ≥24.0'  | ≥45.0'                  | ≥24.0"        |
| 6         | Multi-layered tree | A distinct layer of size class 5 trees over a distinct layer of size class 4 and/or 3 trees, and total tree canopy of the layers ≥60% (layers must have ≥10.0% canopy cover and distinctive height separation). |                         |               |

WHR Codes for Density Classes:

| WHR Code | CWHR Closure Class              | Vegetation Cover (Canopy Closure) |
|----------|---------------------------------|-----------------------------------|
| S        | Sparse Cover                    | 10.0 - 24.9%                      |
| P        | Open Cover                      | 25.0 - 39.9%                      |
| M        | Moderate Cover                  | 40.0 - 59.9%                      |
| D        | Dense Cover                     | ≥60%                              |
| X        | Not Determined / Not Applicable |                                   |

**Data Vintage:** 1990-2023

**Data Resolution:** Raster, 30 meter pixels

**Data Units:** Categorical (see above)

**Creation Method:** Vegetation maps are an important feature of any natural resource management portfolio. Currently the vegetation map for the entire state that is considered the "best available" data is the CALFIRE data known as FVEG (*Vegetation (fveg) - CALFIRE FRAP [ds1327]*). This is an excerpt from the metadata:

"The California Department of Forestry and Fire Protections CALFIRE Fire and Resource Assessment Program (FRAP), in cooperation with California Department of Fish and Wildlife VegCamp program and extensive use of USDA Forest Service Region 5 Remote Sensing Laboratory (RSL) [*now known as Mapping and Remote Sensing*

*Team (MARS)]*, has compiled the "best available" land cover data available for California into a single comprehensive statewide data set. The data span a period from approximately 1990 to 2014. Cross-walks were used to compile the various sources into the common classification scheme, the California Wildlife Habitat Relationships (CWHR) system."

Given the degree of fire in Central Coast in the last 30 plus years, especially in areas that experienced high severity fire, our RRK team thought that using the last version of FVEG (from 2015 but source data could be as old as 1987) would have too many glaring errors. Notwithstanding the challenge of creating reliable vegetation maps, we thought it would be possible to make improvements over the most recent map.

There are many avenues for improving vegetation maps. However, we did not have time to build anything from a new starting point, so we constructed a few simple rules for making [updates to the FVEG data layer](#).

There are three separate rasters provided; one for WHR Veg Type, one for WHR Veg Size Class, and one for WHR Veg Canopy Cover (Density) Class.

The sources for updated data include:

- Fire severity data (from CALFIRE)
- LANDFIRE 2021 land cover data (wildland fire management programs of the USDA Forest Service and USDI)
- Herbaceous cover (Region 5 MARS Team)
- California Forest Observatory (SALO)

**Data Source:** CALFIRE, CDFW, LANDFIRE, California Forest Observatory (SALO), USDA Forest Service

**File Name:** RRK\_Fveg\_WHRtype\_2023Apr\_4regions\_v2.tif; RRK\_Fveg\_WHRsize\_2023Apr\_4regions.tif; RRK\_Fveg\_WHRDensity\_2023Apr\_4regions.tif

## AQUATIC

---

### LAKES/RESERVOIRS

**Definition and Relevance:** Water Bodies such as lake and reservoir features are represented in this layer pulled from the National Hydrography Dataset (NHD). These data were used to erase areas of lakes and ponds from every raster metric in the RRK project dataset.

**Data Resolution:** 30m Raster

**Data Units:** Binary, 0/1

**Creation Method:** This dataset is a subset of vector polygon NHD water bodies, encompassing the RRK project boundary and converted to a raster grid at 30m resolution based on existence/non-existence.

**Data Source:** USGS National Hydrography Dataset (NHD); <https://www.usgs.gov/national-hydrography/national-hydrography-dataset>

**File Name:** NHD\_lakesReservoirs\_2022\_RRK.tif

---

## PERENNIAL, INTERMITTENT AND EPHEMERAL STREAMS

**Definition and Relevance:** USGS National Hydrography Dataset (NHD); Flowline is the fundamental flow network consisting predominantly of stream/river and artificial path vector features. It represents the spatial geometry and carries the attributes

**Data Resolution:** Vector, line

**Data Units:** Tabular attributes

**Creation Method:** Data selected from NHD Flowline feature class to contain only FType code 460, StreamRiver (Perennial, Ephemeral, Intermittent) for the state of California.

**Data Source:** USGS National Hydrography Dataset (NHD); <https://www.usgs.gov/national-hydrography/national-hydrography-dataset>

**File Name:** NHD\_Flowline\_2022\_RRK.tif

## DATA DISCLAIMERS

Appropriate use includes regional assessments of vegetation cover, land cover, or land use change trends, total extent of vegetation cover, land cover, or land use change, and aggregated summaries of vegetation cover, land cover, or land use change. Further use includes applying these data to assess management opportunities for treatments to restore landscape resiliency.

**Attribution** — You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

**ShareAlike** — If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original.

**Non-Commercial Use Only** — Some data layers are restricted by the terms of the data provider. “Non-commercial purposes” means that you may not sell, profit from, or commercialize the content within or works derived from them. Carefully review the terms of each data provider before using the data.

**No additional restrictions** — You may not apply legal terms or technological measures that legally restrict others from doing anything this license permits.

### DEPARTMENT OF FORESTRY AND FIRE PROTECTION (CALFIRE)

The State of California and the Department of Forestry and Fire Protection make no representations or warranties regarding the accuracy of data or maps. The user will not seek to hold the State or the Department liable under any circumstances for any damages with respect to any claim by the user or any third party on account of or arising from the use of data or maps.

### CALIFORNIA DEPARTMENT OF FISH AND WILDLIFE (CDFW)

The state makes no claims, promises, or guarantees about the accuracy, completeness, reliability, or adequacy of these data and expressly disclaims liability for errors and omissions in these data. No warranty of any kind, implied, expressed or statutory, including but not limited to the warranties of non-infringement of third-party rights, title, merchantability, fitness for a particular purpose, and freedom from computer virus, is given with respect to these data.

---

### AREA OF CONSERVATION EMPHASIS (ACE)

The ACE data is subject to certain assumptions and limitations that must be considered in any use or application of the data. All ACE data layers are limited by the accuracy and scale of the input data. ACE is a compilation of the best available scientific information; however, many of these datasets are not comprehensive across the landscape, may change over time, and should be revised and improved as new data become available.

The user accepts sole responsibility for the correct interpretation and use of these data and agrees not to misrepresent these data. CDFW makes no warranty of any kind regarding these data, express or implied. By downloading these datasets, the user understands that these data are in draft condition and subject to change at any time as new information becomes available. The user will not seek to hold the State or the Department liable under any circumstances for any damages with respect to any claim by the user or any third party on account of or arising from the use of data or maps. CDFW reserves the right to modify or replace these datasets without notification.

The ACE maps display biological and recreational values based on available data and constrained by the limitations of the data. The values may be influenced by the level of survey effort in a given area. The ACE data represent broad-scale patterns across the landscape, and the value of any single watershed should be interpreted with caution. ACE is a decision-support tool to be used in conjunction with species-specific information and local-scale conservation prioritization analyses.

The ACE maps do not replace the need for site-specific evaluation of biological resources and should not be used as the sole measure of conservation priority during planning. No statement or dataset shall by itself be considered an official response from a state agency regarding impacts to wildlife resulting from a management action subject to the California Environmental Quality Act (CEQA).

---

## BIOGEOGRAPHIC INFORMATION AND OBSERVATION SYSTEM (BIOS)

Use of this dataset requires prior approval by the primary contact. Recognition that the data set was created and provided by the California Department of Fish and Wildlife, and that any questions regarding the data should be addressed to the contact person listed in the metadata.

### Data Basin - Conservation Biology Institute

Data Basin, by the Conservation Biology Institute (CBI), is a public resource of user-contributed data about conservation issues. Any content including datasets, files, logos, and documents contributed by the user and any resulting data generated by such user belongs to the user, and CBI makes no claim to this content nor does CBI provide any warranty to this content whatsoever. The Data Basin platform itself, and all related documentation, design, and graphic elements (the website as a whole) are the proprietary property of CBI, and CBI possesses all right and title. All of these Data Basin platform rights are reserved.

**Disclaimer** CBI makes no warranty or guarantee as to the content, sequence, accuracy, timeliness, or completeness of any of the information provided herein. CBI explicitly disclaims any representations and warranties of merchantability and fitness for a particular purpose. CBI shall assume no liability for any errors, omissions, or inaccuracies in the information provided regardless of how caused. CBI would appreciate feedback on any errors that are discovered when using this site by contacting us at [databasin@consbio.org](mailto:databasin@consbio.org).

**Use rights** Data Basin has the ability to allow users to browse and search in the Service. You may upload your own data for use within the Data Basin platform; however, this use is limited to only non-commercial purposes. You may not use this site or information found at this site for selling or promoting products or services, soliciting clients, or any other commercial purpose. You may not share your sign-in or password with anyone. You may reproduce, publish, and/or display portions of the Data Basin content only as is necessary to display data for your non-commercial purpose and only if you (1) cite CBI as the owner of Data Basin and (2) abide by any use constraints in citing third-party data. CBI waives any and all liability with respect to unauthorized uses of third-party data. As the user of third-party data, you represent and warrant that you have secured rights in that data and that you will indemnify CBI for any unauthorized use of the data. CBI reserves the sole discretion and right to deny, revoke, or limit use of this site at any time. You may not copy, reproduce, publish, display, make derivative works from, or reverse engineer the Data Basin platform or the Content. You understand and agree that the Service, including all new features provided with the Service, is provided "AS-IS" and that the Provider assumes no responsibility for any content, user communications or personalization settings. You are responsible for obtaining access to the Service and that access may involve third party fees (such as ISP charges). In addition, you must provide and are responsible for all equipment necessary to access the Service.

For additional details see <https://databasin.org/pages/terms-service/>



#### FARMLAND MAPPING AND MONITORING PROGRAM (FMMP)

The State of California and the Department of Conservation make no representations or warranties regarding the accuracy of these data or maps. Neither the State nor the Department shall be liable under any circumstances for direct, indirect, special, incidental or consequential damages with respect to any claim by any user or any third party on account of or arising from the use of these data or maps.

This data does not reflect general plan or zoning designations, city limit lines, changing economic or market conditions, or other factors which may be taken into consideration when land use policies are determined. This data is not designed to be used for parcel specific planning purposes due to its scale and the size of the minimum mapping unit (10 acres).

#### OPEN STREET MAP

This data is made available under the Open Database License: <http://opendatacommons.org/licenses/odbl/1.0/>. Any rights in individual contents of the database are licensed under the Database Contents License: <http://opendatacommons.org/licenses/dbcl/1.0/>. OSM data are free to copy, distribute, transmit and adapt. OpenStreetMap® is open data, licensed under the Open Data Commons Open Database License (ODbL) by the OpenStreetMap Foundation (OSMF).

#### LANDFIRE

LF data are provided "as-is" and without express or implied warranties as to their completeness, accuracy, suitability, or current state thereof for any specific purpose. The LF Program is in no way condoning or endorsing the application of these data for any given purpose. The DOI and USFS manage multiple sets of information and derived data as a service to users of digital geographic data and various databases. No agent of LF shall have liability or responsibility to data users or any other person or entity with respect to any loss or damage caused or alleged to be caused directly or indirectly by the data set. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. government.

#### CALIFORNIA FOREST OBSERVATORY (SALO SCIENCES)

Neither we, the Collaborators, nor our licensors or suppliers make any representations or warranties concerning any content contained in or accessed through the Services, and we will not be responsible or liable for the accuracy, copyright compliance, legality, or decency of material contained in or accessed through the Services. We (and our licensors and suppliers) make no representations or warranties regarding suggestions or recommendations of services or products offered or purchased through the Services. THE SERVICES AND CONTENT ARE PROVIDED ON AN "AS-IS" BASIS, WITHOUT WARRANTIES OR ANY KIND, EITHER EXPRESS OR IMPLIED, INCLUDING, WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, NON-INFRINGEMENT, OR THAT USE OF THE SERVICES WILL BE UNINTERRUPTED OR ERROR-FREE. SOME STATES DO NOT ALLOW LIMITATIONS ON HOW LONG AN IMPLIED WARRANTY LASTS, SO THE ABOVE LIMITATIONS MAY NOT APPLY TO YOU.

#### OPEN DATA COMMONS OPEN DATABASE LICENSE (ODBL)

Open Data Commons is not a law firm and does not provide legal services of any kind.

Open Data Commons has no formal relationship with you. Your receipt of this document does not create any kind of agent-client relationship. Please seek the advice of a suitably qualified legal professional licensed to practice in your jurisdiction before using this document.

No warranties and disclaimer of any damages. This information is provided 'as is', and this site makes no warranties on the information provided. Any damages resulting from its use are disclaimed.

#### CALIFORNIA OFFICE OF ENVIRONMENTAL HEALTH HAZARD ASSESSMENT (OEHHA)

The State makes no claims, promises, or guarantees about the accuracy, completeness, reliability, or adequacy of these data and expressly disclaims liability for errors and omissions in these data. No warranty of any kind, implied, expressed, or statutory, including but not limited to the warranties of non-infringement of third party rights, title, merchantability, fitness for a particular purpose, and freedom from computer virus, is given with respect to these data.

#### CENTER FOR ECOSYSTEM CLIMATE SOLUTIONS (CECS) – UC IRVINE

The University of California ("UC") makes the materials on this website available pursuant to the following disclaimers: the materials are offered "as is"; user assumes any and all risks, of any kind or amount, of using these materials; user shall use the materials only in accordance with law; user releases, waives, discharges and promises not to sue UC, its directors, officers, employees or agents, from liability from any and all claims, including the negligence of UC, resulting in personal injury (including death), accidents or illnesses, property loss, as well as any and all loss of business and/or profit in connection with user's use of the materials; and user shall indemnify and hold UC harmless from any and all claims, actions, suits, procedures, costs, expenses, damages, and liabilities, including attorney's fees, arising out of user's use of the materials and shall reimburse UC for any such incurred expenses, fees or costs.

#### PYROLOGIX

The user must be aware of data conditions and must ultimately bear responsibility for the appropriate use of the information with respect to possible errors, possible omissions, map scale, data collection methodology, data currency, and other conditions specific to certain data.

#### USDA FOREST SERVICE (USFS)

The USDA Forest Service makes no warranty, expressed or implied, including the warranties of merchantability and fitness for a particular purpose, nor assumes any legal liability or responsibility for the accuracy, reliability, completeness, or utility of these geospatial data, or for the improper or incorrect use of these geospatial data. These geospatial data and related maps or graphics are not legal documents and are not intended to be used as such. The data and maps may not be used to determine title, ownership, legal descriptions or boundaries, legal jurisdiction, or restrictions that may be in place on either public or private land. Natural hazards may or may not be depicted on the data and maps, and land users should exercise due caution. The data are dynamic and may change over time. The user is responsible to verify the limitations of the geospatial data and to use the data accordingly.

#### US FISH AND WILDLIFE SERVICE (USFWS)

The use of trade, product, industry or firm names is for informative purposes only and does not constitute an endorsement by the U.S. Fish and Wildlife Service. Link to non-Service Websites do not imply any official U.S. Fish and Wildlife Service endorsement of the opinions or ideas expressed therein or guarantee the validity of the

information provided. Base cartographic information used as part of the Wetlands Mapper has been provided through a license agreement with ESRI and the Department of the Interior.

#### U.S. GEOLOGICAL SURVEY (USGS)

Unless otherwise stated, all data, metadata and related materials are considered to satisfy the quality standards relative to the purpose for which the data were collected. Although these data and associated metadata have been reviewed for accuracy and completeness and approved for release by the U.S. Geological Survey (USGS), no warranty expressed or implied is made regarding the display or utility of the data for other purposes, nor on all computer systems, nor shall the act of distribution constitute any such warranty.

## ADDITIONAL RESOURCES

California Department of Fish and Wildlife Areas of Conservation Emphasis program:

<https://wildlife.ca.gov/Data/Analysis/Ace>

California Department of Fish and Wildlife. California Interagency Wildlife Task Group. 2014. CWHR version 9.0 personal computer program. Sacramento, CA. <http://wildlife.ca.gov/Data/CWHR>

California Office of Environmental Health Hazard Assessment CalEnviroScreen 4.0 report:

<https://oehha.ca.gov/calenviroscreen/report/calenviroscreen-40>

California Forest Observatory (2020). A Statewide Tree-Level Forest Monitoring System. Salo Sciences, Inc. San Francisco, CA. <https://forestobservatory.com>

[Connecting Wildlands & Communities, Conservation Ecology Lab - San Diego State University. Connecting Wildlands & Communities | Climate Science Alliance](#)

Monitoring Trends in Burn Severity (MTBS) program: <https://www.mtbs.gov/>

Multi-Resolution Land Characteristics Consortium (MRLC): <https://www.mrlc.gov/>

Oregon State University Environmental Monitoring, Analysis, and Process Recognition (eMapR) Lab:

<http://emapr.ceoas.oregonstate.edu/>

Rapid Assessment of Vegetation Condition after Wildfire (RAVG): <https://burnseverity.cr.usgs.gov/ravg/>

Spatial Informatics Group: [Home - SIG \(sig-gis.com\)](http://sig-gis.com)

## REFERENCES

Aboud S.A., Spencer L., Wiczorek M., 2022. U.S. Forest Service national riparian areas base map for the conterminous United States in 2019. Fort Collins, CO: Forest Service Research Data Archive.

<https://doi.org/10.2737/RDS-2019-0030>

Bernal, A.A., Stephens, S.L., Collins, B.M., Battles, J.J., 2022. Biomass stocks in California's fire-prone forests: mismatch in ecology and policy. *Environmental Research Letters*. v17, (044047). <https://doi.org/10.1088/1748-9326/ac576a>

Bin Chen and Yufang Jin 2022. Spatial patterns and drivers for wildfire ignitions in California. *Environ. Res. Lett.* 17 055004, DOI 10.1088/1748-9326/ac60da

California Department of Fish and Wildlife. California Interagency Wildlife Task Group. 2014. CWHR version 9.0 personal computer program. Sacramento, CA. <https://wildlife.ca.gov/Data/CWHR>

Carlson, A.R., Helmers, D.P., Hawbaker, T.J., Mockrin, M.H., Radeloff, V.C., 2022. Wildland-urban interface maps for the conterminous U.S. based on 125 million building locations. *Ecological Applications*. v32(5), e2597. <https://doi.org/10.1002/eap.2597> and as U.S. Geological Survey data release: <https://doi.org/10.5066/P94BT6Q7>

Christensen, G.A., Gray, A.N., Kuegler, O., Tase, N.A., Rosenberg, M., Loeffler, D., Anderson, N., Stockmann, K., Morgan, T.A., 2019. *AB 1504 California Forest Ecosystem and Harvested Wood Product Carbon Inventory: 2017 Reporting Period. Final Report*. U.S. Forest Service agreement no. 18-CO-11052021-214, 17-CO-11261979-086, California Department of Forestry and Fire Protection agreement no. 8CA04056 and 8CA03714 and the University of Montana. Sacramento, CA: California Department of Forestry and Fire Protection and California Board of Forestry and Fire Protection. 539 p. <https://www.oregon.gov/ODF/ForestBenefits/Documents/Forest%20Carbon%20Study/Report-CA-1504-forest-ecosys-HWP-CA-2017-13feb19.pdf>

Cowardin, L.M., Carter, V., Golet, F.C., LaRoe, E.T., 1979 Classification of Wetlands and Deepwater Habitats of the United States. U.S. Fish and Wildlife Service, Washington, DC. FWS/OBS 79/31. 103 pp. <https://pubs.er.usgs.gov/publication/2000109>

[Dillon, G. K., Menakis, J., Fay, F. 2015. Wildland fire potential: A tool for assessing wildfire risk and fuels management needs. In: Keane, R. E., Jolly, M., Parsons, R., Riley, K. Proceedings of the large wildland fires conference; May 19-23, 2014; Missoula, MT. Proc. RMRS-P-73. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. p. 60-76.](#)

[Dillon, G. K. 2018. Wildfire Hazard Potential \(WHP\) for the conterminous United States \(270-m GRID\), version 2018 continuous. 2nd Edition. Fort Collins, CO: Forest Service Research Data Archive.](#)

Flint, L.E., Flint, A.L., 2014. California Basin Characterization Model: A Dataset of Historical and Future Hydrologic Response to Climate Change, (ver. 1.1, May 2017): U.S. Geological Survey Data Release, <https://doi.org/10.5066/F76T0JPB>

Girvetz, E. H., Greco, S. E., 2007. How to define a patch: a spatial model for hierarchically delineating organism-specific habitat patches. *Landscape Ecology*. v22, (1131-1142). <https://doi.org/10.1007/s10980-007-9104-8>

Goodwin, M.J., Zald, H.S.J., North, M.P., Hurteau, M.D., 2021. Climate-driven tree mortality and fuel aridity increase wildfire's potential heat flux. *Geophysical Research Letters*. 48, e2021GL094954. <https://doi.org/10.1029/2021GL094954>

Gutierrez, R.J., Manley, P.N., Stine, P.A. tech. eds., 2017. The California spotted owl: current state of knowledge. Gen. Tech. Rep. PSW-GTR-254. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 294 p. [https://www.fs.usda.gov/psw/publications/documents/psw\\_gtr254/](https://www.fs.usda.gov/psw/publications/documents/psw_gtr254/)

Higuera, P. E., Abatzoglou, J. T., 2020. Record-setting climate enabled the extraordinary 2020 fire season in the western United States. *Global Change Biology*. v27(1), (1-2). <https://doi.org/10.1111/gcb.15388>

Jeronimo, S. M. A., Kane, V. R., Churchill, D. J., Lutz, J. A., North, M. P., Asner, G. P., Franklin, J. F., 2019. Forest structure and pattern vary by climate and landform across active-fire landscapes in the montane Sierra Nevada. *Forest Ecology and Management*. v437, (70-86). <https://doi.org/10.1016/j.foreco.2019.01.033>

Keyantash, John & National Center for Atmospheric Research Staff (Eds). Last modified 07 August 2018. "The Climate Data Guide: Standardized Precipitation Index (SPI)." Retrieved 2022 from <https://climatedataguide.ucar.edu/climate-data/standardized-precipitation-index-spi>

Madakumbura, G.D., Goulden, M.L., Hall, A., Fu, R., Moritz, M.A., Koven, C.D., Kueppers, L.M., Norlen, C.A., Randerson, J.T., 2020. Recent California tree mortality portends future increase in drought-driven forest die-off. *Environmental Research Letters*. v15(12), e124040. <https://doi.org/10.1088/1748-9326/abc719>

Mandelbrot, B. B. 1977. *Fractals: Form, Chance, and Dimension*. San Francisco. W. H. Freeman and Company.

Miller, J.D., Knapp, E.E., Key, C.H., Skinner, C.N., Isbell, C.J., Creasy, R.M., Sherlock, J.W., 2009. Calibration and validation of the relative differenced Normalized Burn Ratio (RdNBR) to three measures of fire severity in the Sierra Nevada and Klamath Mountains, California, USA. *Remote Sensing of Environment*. v113(3), (645-656). <https://doi.org/10.1016/j.rse.2008.11.009>

North, M.P., Tompkins, R.E., Bernal, A.A., Collins, B.M., Stephens, S.L., York, R.A., 2022. Operational resilience in western US frequent-fire forests. *Forest Ecology and Management*. v507(12004). <https://doi.org/10.1016/j.foreco.2021.120004>

Parks, S.A., Holsinger, L.M., Voss, M.A., Loehman, R.A., Robinson, N.P., 2018. Mean composite fire severity metrics computed with Google Earth Engine offer improved accuracy and expanding mapping potential. *Remote Sensing*. v10(6), 879. <https://doi.org/10.3390/rs10060879>

Safford, H. D., Van de Water, K. M, 2014. Using fire return interval departure (FRID) analysis to map spatial and temporal changes in fire frequency on national forest lands in California. Res. Pap. PSW-RP-266. [https://www.fs.usda.gov/psw/publications/documents/psw\\_rp266/psw\\_rp266.pdf](https://www.fs.usda.gov/psw/publications/documents/psw_rp266/psw_rp266.pdf)

Scott, J.H., 2020. A deterministic method for generating flame-length probabilities. In: Hood, S.M., Drury, S., Steelman, T., Steffens, R., [eds.]. 2020. Proceedings of the Fire Continuum-Preparing for the future of wildland fire; 2018 May 21-24; Missoula, MT. Proceedings RMRS-P-78. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 358 p. <https://www.fs.usda.gov/treesearch/pubs/62336>

Scott, J. H., Thompson, M. P., Calkin, D. E., 2013. A wildfire risk assessment framework for land and resource management. Gen. Tech. Rep. RMRS-GTR-315. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research station. 83 p. <https://doi.org/10.2737/rmrs-gtr-315> ; <https://www.fs.usda.gov/treesearch/pubs/56265>

Short, Karen C. 2022. Spatial wildfire occurrence data for the United States, 1992-2020 [FPA\_FOD\_20221014]. 6th Edition. Fort Collins, CO: Forest Service Research Data Archive. <https://doi.org/10.2737/RDS-2013-0009.6>

Stephens, S. L., Bernal, A. A., Collins, B. M., Finney, M. A., Lautenberger, C., Saah, D., 2022. Mass fire behavior created by extensive tree mortality and high tree density not predicted by operational fire behavior models in the southern Sierra Nevada. *Forest Ecology and Management*. v518, (120258).  
<https://doi.org/10.1016/j.foreco.2022.120258>

Thorne, J. H., R. M. Boynton, L. E. Flint, and A. L. Flint. 2015. The magnitude and spatial patterns of historical and future hydrologic change in California's watersheds. *Ecosphere* 6(2):24. <http://dx.doi.org/10.1890/ES14-00300.1>

Thorne, J.H., R.M. Boynton, A.J. Holguin, J.A.E. Stewart, & J. Bjorkman. (2016) A climate change vulnerability assessment of California's terrestrial vegetation. California Department of Fish and Wildlife (CDFW), Sacramento, CA.

Thorne, J. H., H. Choe, R. M. Boynton, J. Bjorkman, W. Albright, K. Nydick, A. L. Flint, L. E. Flint, and M. W. Schwartz. 2017. The impact of climate change uncertainty on California's vegetation and adaptation management. *Ecosphere* 8(12):e02021. [10.1002/ecs2.2021](https://doi.org/10.1002/ecs2.2021)

Thorne, J. H., M. Gogul-Prokurat, S. Hill, D. Walsh, R. M. Boynton, H. Choe. 2020. Vegetation refugia can inform climate-adaptive land management under global warming. *Front Ecol Environ* 2020; 18(5): 281–287, doi:10.1002/fee.2208

U.S. Department of Commerce, Census Bureau (USDCCB), 2021. *2020 Census State Redistricting Data (Public Land 94-171) Summary File*, [online resource], <https://www.census.gov/programs-surveys/decennial-census/about/rdo/summary-files.html>

USDA Forest Service. 2018. Toward Shared Stewardship Across Landscapes: An Outcome-Based Investment Strategy. Publication FS-1118. <https://www.fs.usda.gov/sites/default/files/toward-shared-stewardship.pdf>

USDA Forest Service. 2019. Conservation Strategy for the California spotted owl (*Strix occidentalis occidentalis*) in the Sierra Nevada. Publication R5-TP-043. [https://www.fs.usda.gov/Internet/FSE\\_DOCUMENTS/fseprd624135.pdf](https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/fseprd624135.pdf)

U.S. Fish and Wildlife Service. Revised 2019 (2009). A System for Mapping Riparian Areas in The Western United States. Ecological Services Division of Budget and Technical Support Branch of Geospatial Mapping and Technical Support. Falls Church, VA. <https://www.fws.gov/wetlands/documents/A-System-for-Mapping-Riparian-Areas-in-The-Western-United-States-2019.pdf>

Van de Water, K. M., Safford H. D., 2011. A Summary of Fire Frequency Estimates for California Vegetation Before Euro-American Settlement. *Fire Ecology*. v7(3), (26-58).  
<https://fireecology.springeropen.com/track/pdf/10.4996/fireecology.0703026.pdf>

Volger, K.C., Brough, A., Moran, C.J., Scott, J.H., Gilbertson-Day, J.W., 2021. Contemporary Wildfire Hazard Across California. Prepared for: Pacific Southwest Region, USDA Forest Service. Available at: <http://pyrologix.com/reports/Contemporary-Wildfire-Hazard-Across-California.pdf>

Wang J.A., Randerson J. T., Goulden M.L., Knight C.A., & Battles, J.J. (2022). Losses of tree cover in California driven by increasing fire disturbance and climate stress. *AGU Advances*, 3, e2021AV000654.  
<https://doi.org/10.1029/2021AV000654>